



## Vademecum SAVAL

Publication date: 2023-04-12

# Vademecum SAVAL

## Index of products

AB	DRONAVAL PACK	MUXOL	TRAVOF SP
AB ANTITUSIVO	DUALTEN	NEUROVAL CD	TRAVOF T SP
ACTAN / ACTAN CD	DUOVAL	NEURUM	TREX / TREX FORTE
ADAX	ELIXINE	NICODROPS	TRIM 300
AERO-ITAN	ENALTEN	NICOL	TRIOFENO
ALERTEX	ENALTEN-D / ENALTEN-DN	NICOTEARS	TRIOVAL
ALEXIA / ALEXIA FORTE	ETEROVAL	NICOTEARS GEL	TRIOVAL DÍA y NOCHE
ALEXIA-D	EUROCOR	NIRVAN	TRUXIMA
ALIZON	EUROCOR AM	NOVOTEARS	UNDERAN
ALIZON 20	EUROCOR D	NYSKIN	VALAX
ALTAZINC	EUROGESIC	OBEXOL	VALAX D
AMARILLA	EUROGESIC FORTE	OFTABIÓTICO	VALAXAM
AMINTA	EUROGREL	OFTABIÓTICO UNGÜENTO	VALAXAM D
AMOVAL	EUROMICINA	OFTAFILM	VENLAX
AMOVAL 1 GRAMO	EUROMUCIL	OFTAFILM SP	VERTIUM
AMOVAL DUO / 400 / 800 / 1000	EUROVIR	OFTAGEN	VITAYDE-C
ANTALIN / ANTALIN FORTE	EUROVIR CREMA	OFTAGEN COMPUESTO	XOLOF-D
ANTIAX	EVOCAZ CD	OFTALER / OFTALER FORTE	ZAFIN
ARIANA	FIBROLOW LIDOSE	OFTALIRIO	ZIVAL
ATROPINA	FIBROTINA LIDOSE	OFTASONA-N	ZIVAL SOLUCIÓN
BEOF	FIBROX XR	OFTASONA-P	ZOMEL
BEQUIUM	FINEX	OFTAVIR	ZOMEL HP
BIOMOOD	FOSVAL	OFTAVITA	
BLAXITEC	FUNGIUM CREMA	OFTIC	
BLOX	GESIX	OFTOL / OFTOL FORTE	
BLOX-D	GINTOL	OFTOL PLUS	
BRIMOF	HEMOVAL	OLOF	
BRINZOF-T	HIALOF HIDRA	OTICUM	
BRONCATOX	HIDRIUM	PAXON	
BUXON	IDON	PERTIUM	
CALDEVAL	IPSON / IPSON FORTE	PILOCARPINA	
CEFIRAX	IPSON-D / IPSON-D FORTE	PROSTOP-D	
CELTIUM	ITAN	PROTIUM - F	
CIPRODEX	KEVAL	PROTIUM - I	
CIPRODEX OTICO	LATOF	PROTIUM - T	
CIPROVAL	LATOF SP	PRUVAL	
CIPROVAL OFTÁLMICO	LATOF-T	QUETIUM	
CIPROVAL ÓTICO	LATOF-T SP	REALTA	
CLAVINEX	LEOVAL	REMSIMA	
CLAVINEX DUO CD	LEOVAL INYECTABLE	RINOVAL	
CLAVINEX DUO y CLAVINEX DUO FORTE	LEOVAL SOLUCION	RIVOXA 10 mg	
CLIMAVAL	LIFTER	RIVOXA 15 mg y 20 mg	
CLONEX CD	LOMEX IV	RUX	
CLORANFENICOL	LOWDEN	SALCAL	
DEPRAX	LUDIUM 20 MG	SOMNO	
DERMABIOTICO	LUDIUM 5 MG	SOMNO XR	
DERMOSONA	LUKANEX	TELLMI	
DESPEVAL	MENTANIA	TELLMI D	
DIAREN	MIGRAX	TIOF	
DIGENIL	MIGTAL	TIOF PLUS	
DOLODROPS	MOXAVAL	TOL 12 FORTE INYECTABLE	
DOLOVERINA	MOXOF	TOL 12 ORAL / TOL 12 FORTE	
DRONAVAL	MUXELIX	TOL TOTAL JARABE	

## **AB**

### **Bucco-pharyngeal antiseptic**

#### **Composition:**

Each tablet contains:  
Chlorexidine hydrochloride 5 mg  
Excipients q.s.

#### **Presentations:**

12 tablet pack

#### **Indications:**

Mild mouth and throat infections, stomatitis, glossitis, gingivitis and pharyngitis. Dental inflammatory processes; pre and post dental surgery. Antiseptic for mouth and throat, oropharynge, upper buccal vestibule, tonsils, buccal mucous membrane, sublingual region, palate

## **AB ANTITUSIVO**

**Bucco-pharyngeal antiseptic / Antitussive**

### **Composition:**

Each tablet contains:  
Chlorhexidine hydrochloride 5 mg  
Noscapine 10 mg  
Excipients q.s.

### **Presentations:**

12 tablet pack

### **Indications:**

Cough associated with: mild mouth and throat infections, stomatitis, glossitis, gingivitis and pharyngitis. Cough symptomatic treatment, in case of laryngitis, pharyngitis and tracheitis. Mouth and throat antiseptic and antitussive drug for relieving cough which is generally associated with laryngitis, pharyngitis and tracheitis.

## **ACTAN / ACTAN CD**

### **Antidepressant**

#### **Composition:**

Each ACTAN capsule contains:  
Fluoxetine hydrochloride 20 mg  
Excipients q.s.

Each ACTAN CD dispersible tablet contains:  
Fluoxetine hydrochloride 20 mg  
Excipients q.s.

#### **Presentations:**

30 or 60 capsule packs  
30 dispersible scored tablet packs

#### **Indications:**

Fluoxetine hydrochloride is indicated for the acute and maintenance treatment of depression, associated or not with anxiety; for the treatment of obsessive-compulsive disorder in adults and children 8 years and above; for the treatment of premenstrual dysphoric disorder.

It is indicated as an adjuvant to psychotherapy for the reduction of binge-eating and purging activity in patients with bulimia nervosa.

# ADAX

## Antianxiety

### Composition:

Each ADAX tablet contains:

Alprazolam 0,25 mg

Excipients q.s.

Alprazolam 0,50 mg

Excipients q.s.

Alprazolam 1,0 mg

Excipients q.s.

Each ADAX RETARD sustained release tablet contains:

Alprazolam 0,5 mg

Excipients q.s.

Alprazolam 1,0 mg

Excipients q.s.

Alprazolam 2,0 mg

Excipients q.s.

### Presentations:

ADAX: 30 scored tablet pack: 0.25 mg, 0.50 mg and 1 mg

ADAX RETARD: 30 sustained-release tablet pack: 0.5 and 1 mg

### Indications:

Neurotic or reactive depression.

For the management of anxiety disorders as a coadjuvant treatment for anxiety associated with depression.

In the treatment of panic disorders with or without agoraphobia.

## **AERO-ITAN**

**Prokinetic / Antianxiety / Antiflatulent**

### **Composition:**

Each capsule contains:  
Chlordiazepoxide hydrochloride 5 mg  
Metoclopramide hydrochloride 5 mg  
Dimethyl polysiloxane 100 mg  
Excipients q.s.

### **Presentations:**

20 capsule pack

### **Indications:**

AEROITAN is indicated in stress-related dyspeptic conditions presenting with flatulence, postprandial fullness, aerophagy and meteorism. Gastric hypokinesia.

# ALERTEX

## Vaso-dilatant / Brain Stimulant

### Composition:

Each tablet contains:

Modafinil 100 mg

Excipients q.s.

Modafinil 200 mg

Excipients q.s.

### Presentations:

100 and 200 mg Alertex tablets: 30 tablet pack

### Indications:

Modafinil is indicated to improve alertness in patients with excessive daytime sleepiness associated with narcolepsy.

Uses:

Modafinil is indicated to improve wakefulness in patients with excessive sleepiness associated with:

- Narcolepsy: a CNS disorder characterized by somnolence, often accompanied by sudden attacks of weakness (catalepsy), and nighttime sleep alteration. Occasional hypnagogic hallucinations (the state between waking and sleeping) and/or sleep paralysis before falling asleep and waking up. The disorder involves wakefulness and sleep deregulation.
- Obstructive sleep apnea/hypopnea syndrome. (OSAHS)
- Shift work sleep disorder (SWSD).



## **ALEXIA / ALEXIA FORTE**

### **Antihistamine**

#### **Composition:**

Each ALEXIA tablet contains:  
Fexofenadine hydrochloride 120 mg  
Excipients q.s.

Each ALEXIA FORTE tablet contains:  
Fenofexadine hydrochloride 180 mg  
Excipients q.s.

#### **Presentations:**

ALEXIA: 10 and 30 coated tablet packs  
ALEXIA FORTE: 10 and 30 coated tablet packs

#### **Indications:**

Fexofenadine is indicated for the treatment of chronic idiopathic urticaria and seasonal allergic rhinitis.

## **ALEXIA-D**

**Antihistamine / Decongestant**

### **Composition:**

Each capsule contains:  
Fexofenadine hydrochloride 60 mg  
Pseudoephedrine sulfate 120 mg  
Excipients q.s.

### **Presentations:**

10 and 20 capsule packs

### **Indications:**

For the relief of symptoms associated with seasonal allergic rhinitis in adults and children over 12 years of age.

## **ALIZON**

### **Oral Contraceptive**

#### **Composition:**

Each active, pink coated tablet contains:

Chlormadinone acetate 2 mg

Ethinyl estradiol 0.03 mg

Excipients q.s.

Each placebo, white coated tablet contains:

Excipients q.s.

#### **Presentations:**

Pack containing 28 coated tablets

#### **Indications:**

Prevention of pregnancy and is useful for treating signs of androgenization, such as acne.

## **ALIZON 20**

### **Oral Contraceptive**

#### **Composition:**

Each active, pink coated tablet contains:

Chlormadinone acetate 2 mg

Ethinyl estradiol 0.02 mg

Excipients q.s.

Each placebo, white coated tablet contains:

Excipients q.s.

#### **Presentations:**

Pack containing 28 coated tablets

#### **Indications:**

Hormonal contraception. Treatment of moderate acne in women who have been indicated this contraceptive.

# ALTAZINC

## Zinc Supplement

### Composition:

Each ml of oral drops solution contains:

Zinc (as sulfate heptahydrate) 5 mg

Excipients q.s.

Each tablet contains:

Zinc (as sulfate heptahydrate) 10 mg

Excipients q.s.

Zinc (as sulfate heptahydrate) 15 mg

Excipients q.s.

### Presentations:

Bottle with 30 ml oral solution for drops: 5 mg / ml (20 drops)

Pack with 40 score tablets of 10 and 15 mg

### Indications:

Prophylaxis and treatment of zinc deficiency conditions caused by low contribution or poor intestinal absorption, and other conditions that interfere with the use thereof or that increase its body loss, with prior determination of zinc plasma levels.

Uses:

It is used in conditions requiring specific protection such as: child growth, skeletal development, surgical procedure, nutritional deficiency, and zinc deficiency. Coadjuvant in the therapy of hypogonadism and hypothyroidism.

## **AMARILLA**

### **Ophthalmic antiseptic**

#### **Composition:**

Each 100 g of ophthalmic ointment contains:

Mercury yellow oxide 2 g (2%)

Excipients q.s.

#### **Presentations:**

Tube containing 4 g of ophthalmic ointment.

#### **Indications:**

Treatment of minor eye infections.

## **AMINTA**

### **Oral Contraceptive**

#### **Composition:**

Each coated tablet contains:

Desogestrel 75 mcg

Excipients q.s.

#### **Presentations:**

Pack containing 28 coated tablets

#### **Indications:**

Selective oral contraceptive to be used during lactation or when estrogens are contraindicated.

# AMOVAL

## Antibiotic

### Composition:

Each AMOVAL tablet contains:  
Amoxicillin (as trihydrate) 500 mg  
Excipients q.s.

Amoxicillin (as trihydrate) 750 mg  
Excipients q.s.

Each 5 ml of AMOVAL oral suspension contains:  
Amoxicillin (as trihydrate) 250 mg  
Excipients q.s.

Amoxicillin (as trihydrate) 500 mg  
Excipients q.s.

### Presentations:

AMOVAL: 21 tablet pack  
Oral Suspension: Packs with powder to prepare 100ml of suspension

### Indications:

Treatment of infections of the upper and lower respiratory tract, infections of the skin and soft tissues, intra-abdominal sepsis, osteomyelitis, infections of the urinary tract caused by susceptible microorganisms, supported by an antibiogram.



## **AMOVAL 1 GRAMO**

### **Antibiotic**

#### **Composition:**

Each tablet contains:  
Amoxicillin (As trihydrate) 1000 mg  
Excipients q.s.

#### **Presentations:**

14 tablet pack  
20 tablet pack

#### **Indications:**

Treatment of airway bacterial infections. Indicated in infections caused by sensitive germs in localized tissues such as gastrointestinal infections, urinary infections, gonorrhea, skin, soft tissue infections.

Indicated for the eradication of *Helicobacter pylori* in triple therapy with Clarithromycin and omeprazole or esomeprazole.

## **AMOVAL DUO / 400 / 800 / 1000**

### **Antibiotic**

#### **Composition:**

Each 5 ml of reconstituted suspension contains:

Amoxicillin (as trihydrate) 400 mg

Excipients q.s.

Amoxicillin (as trihydrate) 800 mg

Excipients q.s.

Amoxicillin (as trihydrate) 1000 mg

Excipients q.s.

#### **Presentations:**

Amoval Duo 400: packs to reconstitute 70 ml of suspension

Amoval Duo 800: packs to reconstitute 35 and 70 ml of suspension

Amoval Duo 1000: packs to reconstitute 50 and 90 ml of suspension

#### **Indications:**

Indicated in the treatment of airway infections caused by sensitive germs. Indicated also in infections of other localizations caused by sensitive germs, such as otitis media, skin and soft tissue infections, urinary infections, gonorrhoea and gastrointestinal infections.

## **ANTALIN / ANTALIN FORTE**

**Antidepressant / Antianxiety**

### **Composition:**

Each ANTALIN tablet contains:  
Amitriptyline hydrochloride 12,5 mg  
Chlordiazepoxide (as hydrochloride) 5 mg  
Excipients q.s.

Each ANTALIN FORTE tablet contains:  
Amitriptyline hydrochloride 25 mg  
Chlordiazepoxide (as hydrochloride) 10 mg  
Excipients q.s.

### **Presentations:**

30 tablet packs of ANTALIN or ANTALIN Forte

### **Indications:**

ANTALIN and ANTALIN FORTE are indicated for the treatment of mild to moderate depressive disorders; accompanied by mild to moderate anxiety episodes.

## **ANTIAX**

**Antacid / Antiflatulent**

### **Composition:**

Each tablet contains:

Magaldrate 480 mg

Simethicone 100 mg

Excipients q.s.

Each 5 ml of oral suspension contains:

Magaldrate 480 mg

Simethicone 100 mg

Excipients q.s

### **Presentations:**

6 and 24 tablet packs

180 and 360 ml suspension bottles

### **Indications:**

Treatment of dyspepsia, flatulence, and heartburn

# ARIANA

## Oral Contraceptive

### Composition:

Each active, white coated tablet contains:

Nomegestrol Acetate 2.5 mg

Estradiol 1.5 mg

Excipientes: q.s.

Each placebo, red coated tablet contains:

Excipients q.s.

### Presentations:

Pack containing 28 coated tablets

### Indications:

Oral contraception.

The decision to prescribe Ariana must take into account the current risk factors of the woman in particular, specifically those of venous thromboembolism, and how the risk of venous thromboembolism with Ariana compares with that of other combined hormonal contraceptives.

# ATROPINA

## Mydriatic-Cycloplegic

### Composition:

Each ml of sterile ophthalmic solution contains:

Atropine sulfate 10 mg (1%)

Excipients q.s.

Atropine sulfate 5 mg (0,5%)

Excipients q.s.

### Presentations:

Sealed bottle containing 10 ml of sterile ophthalmic solution

### Indications:

The ophthalmic preparations of Atropina are used to produce mydriasis and cyclopegia during retinal and optic disc examination, together with the refractive error measurement.

Atropina is also used in the management of acute inflammatory processes of the iris (iridocyclitis), and the uveal tract (uveitis). Atropina is also used for its cyclopegic effect for the treatment of amblyopia suppression by reducing visual acuity in the non-affected eye, while the affected eye forces fixation. It has also been used to treat patients with a functional excess of accommodation and convergence.

## **BEOF**

### **Antiglaucoma**

#### **Composition:**

Each ml of sterile ophthalmic solution contains:

Betaxolol hydrochloride 5 mg (0,5%)

Excipients q.s.

#### **Presentations:**

Sealed drop dispenser bottle with 5 ml of sterile ophthalmic solution

#### **Indications:**

BEOF is indicated for the treatment of ocular hypertension and open angle chronic glaucoma. It may be used alone or combined with other medications for glaucoma.

## **BEQUIUM**

**Antitussive / Antihistamine / Decongestant**

### **Composition:**

Each 5 ml of oral solution contains:  
Chlorphenamine maleate 2 mg  
Pseudoephedrine hydrochloride 30 mg  
Codeine 10 mg  
Excipients q.s.

### **Presentations:**

120 ml bottle

### **Indications:**

BEQUIUM is indicated for the symptomatic treatment of non-productive cough, allergic, irritative in origin, associated with bronchitis and/or cold involving nasal congestion.



# BIOMOOD

## Antiepileptic

### Composition:

Each coated tablet contains:  
Topiramate 25, 50 or 100 mg  
Excipients q.s.

### Presentations:

BIOMOOD® 25 mg: 30 coated tablets pack  
BIOMOOD® 50 mg: 30 coated tablets pack  
BIOMOOD® 100 mg: 30 coated tablets pack

### Indications:

BIOMOOD® is indicated in:

- Epilepsy
  - Indicated as initial monotherapy in patients 6 years of age and older with partial onset or primary generalized tonic-clonic seizures
  - Indicated as an add-on therapy in epileptic adult patient and children between 2 and 16 years having focal seizures or focal onset which then generalize
  - Indicated in adults and children over 2 years as an add-on therapy of crisis associated with Lennox-Gastaut syndrome
- Migraine
  - Indicated in adult patients for prophylaxis of migraine

The acute treatment of migraine has not been studied

## **BLAXITEC**

**Antihistamine**

### **Composition:**

Each tablet contains 20 mg of bilastine.

### **Presentations:**

Package with 30 tablets

### **Indications:**

Symptomatic treatment of allergic rhinoconjunctivitis (seasonal and perennial) and urticaria. Blaxitec 20 mg tablets are indicated in adults and adolescents (age equal to or greater than 12 years).

# **BLOX**

## **Antihypertensive**

### **Composition:**

Each tablet contains:

Candesartan cilexetil 8 mg

Excipients q.s.

Candesartan cilexetil 16 mg

Excipients q.s.

Candesartan cilexetil 32 mg

Excipients q.s.

### **Presentations:**

Blox 8 mg: package with 30 tablets

Blox 16 mg: package with 30 tablets

Blox 32 mg: package with 30 tablets

### **Indications:**

It is indicated for:

Treatment of High Blood Pressure.

Treatment of patients with heart failure and left ventricular systolic dysfunction (ejection fraction less than or equal to 40%), as a complement of ACE inhibitors, or in case of intolerance to this kind of medications.

## **BLOX-D**

**Antihypertensive / Diuretic**

### **Composition:**

Each tablet contains:

Candesartan cilexetil 8 mg

Hydrochlorothiazide 12,5 mg

Excipients q.s.

Candesartan cilexetil 16 mg

Hydrochlorothiazide 12,5 mg

Excipients q.s.

Candesartan cilexetil 32 mg

Hydrochlorothiazide 12,5 mg

Excipients q.s.

### **Presentations:**

30 scored tablet packs of:

BLOX-D 8/12.5

BLOX-D 16/12.5

BLOX-D 32/ 12.5

### **Indications:**

This product is indicated for the treatment of idiopathic arterial hypertension in patients not achieving an appropriate control thereof, with a monotherapy with candesartan cilexethyl or hydrochlorothiazide.

## **BRIMOF**

### **Antiglaucoma**

#### **Composition:**

Each 1 mL (28 drops) of ophthalmic solution contains:

Brimonidine Tartrate 2 mg

Excipients: q.s.

Each 100 ml of ophthalmic solution contains:

Brimonidine Tartrate 0,2 g

Excipients q.s.

#### **Presentations:**

Sealed drop bottle with 10 ml of sterile ophthalmic solution

#### **Indications:**

BRIMOF is indicated for the reduction of elevated intraocular pressure in patients with open-angle glaucoma and ocular hypertension.

## **BRINZOF-T**

### **Antiglaucoma**

#### **Composition:**

Each 1 mL (27 drops) of ophthalmic suspension contains:

Brinzolamide 10mg  
Timolol (as maleate) 5 mg  
Excipients q.s.

Each 100 mL of ophthalmic suspension contains:

Brinzolamide 1.0 g  
Timolol (as maleate) 0.5 g  
Excipients q.s.

#### **Presentations:**

Sealed drop bottle with 5 mL of sterile ophthalmic suspension

#### **Indications:**

Lowering of intraocular pressure (IOP) in adult patients with open-angle glaucoma or ocular hypertension for whom monotherapy does not provide sufficient IOP reduction.

# BRONCATOX

**Antitussive**

## Composition:

Each 10 ml of oral solution contains:

Levodropropizine 60 mg

Excipients q.s.

## Presentations:

120 ml bottle

## Indications:

Symptomatic treatment of non-productive cough

## **BUXON**

### **Antidepressant / Smoking Cessation Agent**

#### **Composition:**

Each tablet contains:  
Anfebutamona hydrochloride (Bupropion) 150 mg  
Excipients q.s.

#### **Presentations:**

Aluminum blister/aluminum packs containing 30 and 60 coated sustained-release tablets

#### **Indications:**

Indicated for the treatment of depression.  
Indicated as a coadjuvant in the treatment of smoking cessation.



## **CALDEVAL**

### **Vitamins and minerals supplement**

#### **Composition:**

Each chewable tablet contains:  
Calcium carbonate 1250 mg  
(Equivalent to 50 mg of ion calcium)  
Vitamin D3 400 I.U  
Excipients q.s.

#### **Presentations:**

Blister pack with 30 and 60 pleasant-tasting flavor, scored tablets

#### **Indications:**

Treatment and prophylaxis of calcium deficiency conditions associated with vitamin D deficit. Indicated as stimulant of fracture consolidation processes.

Uses:

Osteoporosis, osteomalacia, rickets, in all forms thereof, hypocalcemia associated with hypoparathyroidism. Breastfeeding, pregnancy, pre- and post-menopausal demineralization and old age.

# CEFIRAX

## Antibiotic

### Composition:

Each tablet contains:  
Cefpodoxime 200 mg  
Excipients q.s.

Each 5 mL of oral suspension contains:  
Cefpodoxime 100 mg  
Excipients q.s.

### Presentations:

Tablets: 10 and 20 tablet pack  
Suspension: Bottle containing 75 mL of oral suspension

### Indications:

Cefpodoxime proxetil is indicated in the oral treatment of respiratory tract infections caused by susceptible microorganisms.

- Acute exacerbations of chronic bronchitis.
- Community-acquired pneumonia.
- Acute maxillary sinusitis.
- Acute otitis media.
- Pharyngitis and tonsillitis caused by streptococcus pyogenes.

It is also indicated in infections caused by susceptible microorganisms:

- Non-complicated gonorrhea.
- Non-complicated skin or skin structure infections.
- Non-complicated urinary tract infections.

# CELTIUM

## Antidepressant

### Composition:

Each coated tablet contains:

Escitalopram (as oxalate) 10 mg

Excipients q.s.

Escitalopram (as oxalate) 20 mg

Excipients q.s.

### Presentations:

30 tablet packs of 10 or 20 mg

60 tablet packs of 10 mg

### Indications:

CELTIUM contains escitalopram, an antidepressant, indicated for:

- Acute treatment and maintenance of major depressive disorder in adults and adolescents 12 years of age.
- Treatment of panic attacks with or without agoraphobia.
- Treatment of social anxiety disorders (social phobia).
- Treatment of generalized anxiety disorders.
- Treatment of obsessive-compulsive disorders.

# CIPRODEX

## Ophthalmic Antibiotic / Corticosteroid

### Composition:

Each 100 g of ophthalmic ointment contains:  
Ciprofloxacin (as hydrochloride monohydrate) 0,3 g  
Dexamethasone 0,1 g  
Excipients q.s.

Each 100 ml of ophthalmic suspension contains:  
Ciprofloxacin (as hydrochloride monohydrate) 0,3 g  
Dexamethasone 0,1 g  
Excipients q.s.

### Presentations:

Ophthalmic suspension: sealed eye drop bottle with 5 mL of sterile ophthalmic suspension  
Ophthalmic ointment: ointment tubes with 3.5 g

### Indications:

Ocular infections caused by susceptible microorganisms when the anti-inflammatory action of dexamethasone is needed. Blepharitis, blepharoconjunctivitis and conjunctivitis caused by sensitive germs, including *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Streptococcus pneumoniae*. Prophylaxis in ocular surgery and post-surgery.

# CIPRODEX OTICO

**Otic Antibiotic / Corticosteroid**

## Composition:

Each ml of otic suspension contains:  
Ciprofloxacin (as hydrochloride monohydrate) 3 mg  
Dexamethasone 1 mg  
Excipients q.s.

## Presentations:

7.5 ml sterile suspension bottle

## Indications:

Otic ciprodex is indicated in:

- Acute otitis media with tympanostomy due to an infection caused by microorganisms sensitive to Ciprofloxacin children older than 6 months.
- Acute external otitis due to an infection caused by microorganisms sensitive to Ciprofloxacin.

# CIPROVAL

## Antibiotic

### Composition:

Each CIPROVAL tablet contains:  
Ciprofloxacin (as hydrochloride monohydrate) 250 mg  
Excipients q.s.

Ciprofloxacin (as hydrochloride monohydrate) 500 mg  
Excipients q.s.

### Presentations:

Pack with 14 tablets of 250 mg  
Pack with 10 and 20 tablets of 500 mg

### Indications:

This medication is for the treatment of single or mixed infections caused by two or more sensitive microorganisms, for example, respiratory tract infections, urinary infections, gastrointestinal infections, gonorrhoea. Additionally, it is an alternative agent against infections caused by antibiotic resistance mechanisms.

# CIPROVAL OFTÁLMICO

## Ophthalmic Antibiotic

### Composition:

Each 100 ml of sterile ophthalmic solution contains:  
Ciprofloxacin (as hydrochloride) 0,3 g  
Excipients q.s.

Each 100 g of sterile ophthalmic ointment contains:  
Ciprofloxacin (as hydrochloride) 0,3 g  
Excipients q.s.

### Presentations:

Sealed bottle with 5 ml sterile ophthalmic solution.  
Ointment container with 3.5 g sterile ophthalmic ointment.

### Indications:

Clinical indications for the oftalmic solution:

CIPROVAL is indicated for the treatment of bacterial conjunctivitis, blepharitis, blepharoconjunctivitis, and corneal ulcer caused by susceptible microorganisms.

Clinical indications for the ophthalmic ointment:

Ophthalmic ciprofloxacin is indicated as an alternative for the treatment of blepharitis and blepharoconjunctivitis caused by susceptible strains.

# CIPROVAL ÓTICO

## Otic Antibiotic

### Composition:

Each 100 ml of sterile solution for ear drops contains:

Ciprofloxacin (as hydrochloride) 0,3 g

Excipients q.s.

### Presentations:

Pack with 5 ml plastic bottle.

### Indications:

CIPROVAL Otic Solution is an antimicrobial quinolone indicated in the treatment of otic infections caused by germs sensitive to ciprofloxacin.



# CLAVINEX

## Antibiotic

### Composition:

Each tablet contains:

Clavulanic acid (as potassium clavulanate) 125 mg

Amoxicillin (as trihydrate) 500 mg

Excipients q.s.

Each 5 ml of oral suspension contains:

Clavulanic acid (as potassium clavulanate) 62,5 mg

Amoxicillin (as trihydrate) 250 mg

Excipients q.s.

### Presentations:

Tablets: 21 tablet pack

Suspension: bottle with powder to prepare 60 ml of 250 mg/5ml suspension

### Indications:

Treatment of infections of the upper and lower respiratory tract, urinary tract, skin and soft tissue, intraabdominal sepsis, osteomyelitis, caused by sensitive microorganisms y beta-lactamase producers supported by an antibiogram.

Clinical uses:

- Acute bacterial sinusitis (properly diagnosed)
- Acute otitis media, tonsillitis
- Acute exacerbation of chronic bronchitis (properly diagnosed)
- Community acquired pneumonia
- Cystitis, urethritis
- Pyelonephritis
- Skin and soft tissue infections, particularly cellulitis, animal bites, severe dental abscesses with spreading cellulitis.
- Bone and joint infections, particularly osteomyelitis.

## **CLAVINEX DUO CD**

### **Antibiotic**

#### **Composition:**

Each dispersible tablet contains:

Amoxicillin 875 mg

Clavulanic acid 125 mg

Excipients q.s.

#### **Presentations:**

14 and 20 dispersible tablets pack

#### **Indications:**

CLAVINEX DUO CD is indicated for the treatment of Beta-lactamase-producing bacteria infections, of the respiratory tract (including ear and nose), urinary tract, skin and soft tissues, intraabdominal sepsis and osteomyelitis caused by organisms sensitive to the association.

# CLAVINEX DUO y CLAVINEX DUO FORTE

## Antibiotic

### Composition:

Each CLAVINEX DUO tablet contains:

Amoxicillin (as trihydrate) 875 mg

Clavulanic acid (as potassium clavulanate) 125 mg

Excipients q.s.

Each 5 ml of CLAVINEX DUO reconstituted suspension contains:

Amoxicillin (as trihydrate) 400 mg

Clavulanic acid (as potassium clavulanate) 57 mg

Excipients q.s.

Each 5 ml of CLAVINEX DUO FORTE reconstituted suspension contains:

Amoxicillin (as trihydrate) 800 mg

Clavulanic acid (as potassium clavulanate) 57 mg

Excipients q.s.

### Presentations:

CLAVINEX Duo

14 tablet and 20 tablet Packs.

Pack containing 1 bottle with powder to reconstitute 35 ml or 70 ml suspension plus 1 bottle with solvent and dosing syringe

CLAVINEX Duo FORTE

Pack containing 1 bottle with powder to reconstitute 35 ml or 70 ml suspension plus 1 bottle with solvent and dosing syringe

### Indications:

Treatment of infections of the upper and lower respiratory tract, urinary tract, skin and soft tissue, intraabdominal sepsis, osteomyelitis, caused by sensitive microorganisms y beta-lactamase producers supported by an antibiogram.

Used in infections caused by microorganisms sensitive to Amoxicillin and infections caused by microorganisms sensitive to Amoxicillin and beta-lactamase producers such as:

- Infections of the upper respiratory tract: e.g.: tonsillitis, sinusitis, and otitis media.
- Infections of the lower respiratory tract: e.g.: acute chronic bronchitis with exacerbations, bronchopneumonia.
- Infections of the genitourinary tract: cystitis, urethritis, pyelonephritis, female genital infections.
- Skin and soft tissue infections.
- Bone and joint infections: e.g. osteomyelitis
- Other infections: e.g.: septic abortion, puerperal sepsis, intraabdominal sepsis, septicemia, peritonitis, post-operative infections.
- Prophylaxis against infections associated with major surgical procedures such as gastrointestinal, pelvis, head and neck, heart, kidney, bone replacement, and biliary tract surgeries.

# CLIMAVAL

## Hormone replacement therapy

### Composition:

Each tablet contains:

Tibolone 2,5 mg

Excipients q.s.

### Presentations:

Pack containing 30 tablets

### Indications:

Climaval is indicated in:

- Treatment of estrogen deficiency symptoms in postmenopausal women, more than a year after menopause.
- Prevention of osteoporosis in postmenopausal women at high risk of fractures in the future who cannot tolerate or have contraindications for other approved medicinal products for the prevention of osteoporosis.

### NOTE

For all women, the decision to prescribe Climaval should be based on an assessment of the general risks of each patient and, in particular in those over 60 years of age, should include consideration of the risk of stroke.

## **CLONEX CD**

**Antianxiety / Anticonvulsivant**

### **Composition:**

Each dispersable tablet contains:

Clonazepam 0,5 mg

Excipients q.s.

Clonazepam 1,0 mg

Excipients q.s.

Clonazepam 2,0 mg

Excipients q.s.

### **Presentations:**

CLONEX 0.5 mg, 1.0 mg and 2.0 mg in 30 dispersible tablet packs

### **Indications:**

It is indicated in clinical epilepsy alone or as adjuvant in the treatment of petit mal seizure (absence) and Lennox-Gastaut syndrome. Also used in akinetic and atonic seizures, mioclonic seizure, primary or secondary generalized tonic-clonic seizure, panic attack and panic disorders with or without agoraphobia (fear to be alone in open spaces and public areas).

# CLORANFENICOL

## Ophthalmic Antibiotic

### Composition:

Each ml of ophthalmic solution contains:  
Chloranphenicol 5 mg  
Excipients q.s.

Each g of ophthalmic ointment contains:  
Chloranphenicol 10 mg  
Excipients q.s.

### Presentations:

10 ml sealed eye drop applicator with sterile ophthalmic solution  
3.5 g ointment tube with ophthalmic ointment

### Indications:

Superficial eye infections such as: acute microbial conjunctivitis, microbial keratitis of lacrimal ducts, blepharitis or other infections caused by sensitive bacteria.

# DEPRAX

## Antidepressant

### Composition:

Each film coated tablet contains:  
Sertraline hydrochloride 50 mg  
Excipients q.s.

Each film coated tablet contains:  
Sertraline hydrochloride 100 mg  
Excipients q.s.

### Presentations:

30 and 60 coated scored tablet pack of Deprax 50 mg  
30 coated scored tablet pack of Deprax 100 mg

### Indications:

Sertraline is indicated for the treatment of depression, including major depressive disorder and relapse prevention, obsessive-compulsive disorder, in adults and pediatric patients aged 6 to 17 years, panic disorder with or without agoraphobia, posttraumatic stress disorder, premenstrual dysphoric disorder, and for the treatment of social anxiety disorder.

## **DERMABIOTICO**

**Antiseptic external use**

### **Composition:**

Each g of ointment contains:  
Polymyxin B (as sulfate) 10000 I.U.  
Bacitracin (as Bacitracin zinc) 500 I.U.  
Excipients q.s.

### **Presentations:**

Topical ointment, 15 g tube.

### **Indications:**

Antibiotic combination for external use. It is used to prevent infections in cuts, abrasions, scratches and minor burns, bacterial dermatoses sensitive to antibiotics, and it is an essential component in first aid kit.



## **DERMOSONA**

### **Topical Corticosteroid**

#### **Composition:**

Each g of cream contains:

Mometasone furoate 1 mg (0,1%)

Excipients q.s.

#### **Presentations:**

Tubes of 10 and 15 g.

#### **Indications:**

Used to relieve inflammatory and itching manifestations of corticosteroid-sensitive dermatoses, such as atopic dermatitis, allergic contact dermatitis and psoriasis.

# DESPEVAL

## Antihistamine

### Composition:

Each coated tablet contains:

Desloratadine 5 mg

Excipients q.s.

Each 5 ml of oral solution contains:

Desloratadine 2,5 mg

Excipients q.s.

### Presentations:

10, 30 and 40 coated tablet packs.

100 ml syrup bottle.

### Indications:

For the relief of symptoms associated with allergic rhinitis. Relief of symptoms of urticaria such as the relief of itching and reduction of wheals size.

# DIAREN

## Antidiarrheal

### Composition:

Each tablet contains:  
Nifuroxazide 200 mg  
Activated attalugite 350 mg  
Excipients q.s.

Each 5 ml of oral suspension contains:  
Nifuroxazide 200 mg  
Activated attalugite 500 mg  
Excipients q.s.

### Presentations:

Tablets: 20 tablet packs  
Suspension: 100 ml bottle

### Indications:

DIAREN is used for acute bacterial diarrheas in children and adults, and chronic diarrheas.

## **DIGENIL**

**Antiflatulent / Digestant**

### **Composition:**

Each capsule contains:

Pancreatin x 6NF 230 mg

(Equivalent to: 5500 FIP U. Amylase, 6500 FIP U. Lipase, 400 FIP U. Protease)

Simethicone 80 mg

Excipients q.s.

### **Presentations:**

Presentation: 10 capsules packs

### **Indications:**

Symptomatic relief of digestive disorders characterized by gastrointestinal flatulence.

## **DOLODROPS**

**Nonsteroidal anti-inflammatory**

### **Composition:**

Each ml (25 drops) contains:  
Diclofenac resinate 46,44 mg  
Excipients q.s.

### **Presentations:**

25 ml bottle

### **Indications:**

Inflammatory rheumatism in children. Post-operative and post-traumatic painful and inflammatory conditions. Adjuvant in severe painful infections of the ear, nose and throat infections.

# DOLOVERINA

## Antispasmodic-Musculotropic

### Composition:

Each sustained-release tablet contains:  
Mebeverine Hydrochloride: 200 mg  
Excipients q.s.

### Presentations:

20 sustained-release tablet pack

### Indications:

Symptomatic treatment of pain and intestinal discomfort due to functional alterations in the gastrointestinal tract and the bile duct.

Clinical uses:

Irritable bowel syndrome, persistent abdominal pain and cramping, non specific diarrhea (with or without alternating constipation), feeling of fullness in the stomach. Treatment of gastrointestinal spasms secondary to organic disorders. Enteritis.

## **DRONAVAL**

**Antiresortive**

### **Composition:**

Each coated tablet contains:

150 mg Ibandronic Acid (as Ibandronate monosodium monohydrate)

Excipients c.s.

### **Presentations:**

3 and 1 tablet packs

### **Indications:**

It is indicated for the treatment and prevention of postmenopausal osteoporosis

## **DRONAVAL PACK**

### **Antiresortive**

#### **Composition:**

Each coated tablet contains:

Ibandronic Acid (as Ibandronate monosodium monohydrate) 150 mg

Excipients q.s.

Each chewable tablet contains:

Calcium carbonate 1250 mg

(Equivalent to 500 mg of ion calcium)

Vitamin D3 400 I.U

Excipients q.s.

#### **Presentations:**

Pack containing: 1 coated tablet of 150 mg Ibandronic Acid and 60 chewable tablets of 500 mg calcium + 400 IU Vitamin D3

#### **Indications:**

It is indicated for the treatment and prevention of osteoporosis

Treatment and prevention of postmenopausal osteoporosis



## DUALTEN

### Antihypertensive

#### Composition:

Each tablet contains:

Carvedilol 6,25 mg

Excipients q.s.

Carvedilol 12,5 mg

Excipients q.s.

Carvedilol 25 mg

Excipients q.s.

#### Presentations:

Dualten 6,25 mg: 30 tablet packs

Dualten 12,5 mg: 30 tablet packs

Dualten 25 mg: 30 tablet packs

#### Indications:

Treatment of the essential Hypertension. Treatment of stable angina. Treatment of symptomatic congestive heart failure.

## **DUOVAL**

### **Antihistamine / Decongestant**

#### **Composition:**

Each ml (24 drops) of drop solution contains:

Chlorphenamine maleate 1 mg  
Pseudoephedrine hydrochloride 30 mg  
Excipients q.s.

Each 5 ml of oral solution contains:

Chlorphenamine maleate 2 mg  
Pseudoephedrine hydrochloride 30 mg  
Excipients q.s.

#### **Presentations:**

Syrup: plastic bottle containing 120 ml solution

Drops: plastic bottle containing 15 ml solution

#### **Indications:**

Congestion of the upper respiratory tract (including the paranasal sinuses and the Eustachian tubeS), common cold, hay fever, vasomotor and allergic rhinitis.

# ELIXINE

## Antiasthmatic

### Composition:

Each 15 ml of oral solution contains:

Theophylline 80 mg

Excipients q.s.

Each lentocaps (sustained release capsule) contains:

Theophylline 125 mg

Excipients q.s.

Theophylline 250 mg

Excipients q.s.

Theophylline 300 mg

Excipients q.s.

### Presentations:

Oral solution: 250 ml bottle

Pack containing 20 lentocaps of 125 mg

Pack containing 20 lentocaps of 250 mg

Pack containing 20 lentocaps of 300 mg

### Indications:

Indicated as bronchodilator in the symptomatic treatment of asthma and reversible bronchospasm associated with chronic bronchitis and emphysema. Indicated for the symptomatic treatment of chronic obstructive pulmonary disease.

# ENALTEN

## Antihypertensive

### Composition:

Each tablet contains:  
Enalapril maleate 5 mg  
Excipients q.s.

Enalapril maleate 10 mg  
Excipients q.s.

Enalapril maleate 20 mg  
Excipients q.s.

### Presentations:

Pack containing 30 scored tablets of 5 mg, 10 mg and 20 mg, in a double laminated hermetic blister

### Indications:

ENALTEN is indicated for the treatment of arterial hypertension and for the treatment of congestive heart failure, concomitantly with other treatments.

## **ENALTEN-D / ENALTEN-DN**

**Antihypertensive / Diuretic**

### **Composition:**

Each ENALTEN D tablet contains:

Enalapril maleate 10 mg  
Hydrochlorotiazide 25 mg  
Excipients q.s.

Each ENALTEN DN tablet contains:

Enalapril maleate 20 mg  
Hydrochlorotiazide 12,5 mg  
Excipients q.s.

### **Presentations:**

ENALTEN-DN 20/12.5: Double laminated hermetic blister pack containing 30 scored tablets

ENALTEN-D (10/25): Double laminated hermetic blister pack containing 30 scored tablets

### **Indications:**

ENALTEN D / ENALTER DN is indicated for the treatment of essential arterial hypertension, resistant to monotherapy.

# ETEROVAL

## Analgesic / Anti-Inflammatory NSAID

### Composition:

Each coated tablet of Eteroval 60 mg contains:

Etoricoxib 60 mg

Excipients q.s.

Each coated tablet of Eteroval 90 mg contains:

Etoricoxib 90 mg

Excipients q.s.

Each coated tablet of Eteroval 120 mg contains:

Etoricoxib 120 mg

Excipients q.s.

### Presentations:

Eteroval 60 mg: package with 14 coated tablets

Eteroval 90 mg: package with 14 coated tablets

Eteroval 120 mg: package with 7 coated tablets

### Indications:

ETEROVAL is indicated in adults and adolescents over 16 years of age for:

Eteroval 60 mg:

- Symptomatic relief of osteoarthritis and rheumatoid arthritis (RA)
- Treatment of ankylosing spondylitis (AS)
- Treatment of primary dysmenorrhea
- Short-term treatment of moderate pain associated with dental surgery
- Pain and signs of inflammation associated with acute gouty arthritis

Eteroval 90 mg:

- Symptomatic relief of rheumatoid arthritis (RA)
- Treatment of ankylosing spondylitis (AS)
- Short-term treatment of moderate pain associated with dental surgery
- Moderate to severe treatment of acute postoperative pain associated with abdominal gynecological surgery

Eteroval 120 mg:

- Treatment of acute gouty arthritis
- Relief of acute pain
- Treatment of primary dysmenorrhea

The decision to prescribe a selective COX-2 inhibitor should be based on an individual assessment of the patient's overall risks.

# EUROCOR

## Antihypertensive

### Composition:

Each coated tablet of Eurocor 1,25 mg contains:

Bisoprolol Fumarate 1,25 mg

Excipients q.s.

Each coated tablet of Eurocor 2,5 mg contains:

Bisoprolol Fumarate 2,5 mg

Excipients q.s.

Each coated tablet of Eurocor 5 mg contains:

Bisoprolol Fumarate 5 mg

Excipients q.s.

Each coated tablet of Eurocor 10 mg contains:

Bisoprolol Fumarate 10 mg

Excipients q.s.

### Presentations:

EUROCOR 1,25 mg: 14 tablet pack

EUROCOR 2,5 mg: 35 tablet pack

EUROCOR 5 mg: 35 tablet pack

EUROCOR 10 mg: 35 tablet pack

### Indications:

Bisoprolol fumarate is indicated in:

- Treatment of arterial hypertension
- Treatment of coronary heart disease
- Treatment of stable chronic heart failure concomitantly with ACE inhibitors, diuretics, and optionally cardiac glycosides

# EUROCOR AM

## Antihypertensive

### Composition:

Each coated tablet of EUROCOR AM 5/5 mg contains:

Bisoprolol Fumarate 5 mg

Amlodipine (as besylate) 5 mg

Excipients q.s.

Each coated tablet of EUROCOR AM 5/10 mg contains:

Bisoprolol Fumarate 5 mg

Amlodipine (as besylate) 10 mg

Excipients q.s.

### Presentations:

EUROCOR AM 5/5 mg: 35 coated tablets pack

EUROCOR AM 5/10 mg: 35 coated tablets pack

### Indications:

EUROCOR AM is indicated in the treatment of arterial hypertension, as a substitution therapy, in patients properly controlled with the individual products administered concomitantly at the same dose level as in the combination, but as separate tablets.



## **EUROCOR D**

**Antihypertensive / Diuretic**

### **Composition:**

Each coated tablet of Eurocor-D 2,5/6,25 contains:

Bisoprolol Fumarate 2,5 mg

Hydrochlorotiazide 6,25 mg

Excipients q.s.

Each coated tablet of Eurocor-D 5/6,25 contains:

Bisoprolol Fumarate 5 mg

Hydrochlorotiazide 6,25 mg

Excipients q.s.

### **Presentations:**

Eurocor D 2,5/6,25: 35 coated tablet pack

Eurocor D 5/6,25: 35 coated tablet pack

### **Indications:**

Eurocor-D (Bisoprolol Fumarate and hydrochlorotiazide) is indicated for the treatment of High Blood Pressure

# EUROGESIC

**Analgesic / Antipyretic / Anti-Inflammatory NSAID / Decongestant**

## Composition:

Each EUROGESIC ADULTO tablet contains:

Naproxen sodium 275 mg

Excipients q.s.

Each EUROGESIC INFANTIL tablet contains:

Naproxen sodium 100 mg

Excipients q.s.

Each EUROGESIC INFANTIL suppository contains:

Naproxen sodium 50 mg

Excipients q.s.

Each ml of EUROGESIC INFANTIL oral suspension contains:

Naproxen sodium 25 mg

Excipients q.s.

## Presentations:

Pack containing 10 adult tablets

Pack containing 10 pediatric tablets

Pack containing 6 suppositories

60 ml oral suspension bottle

## Indications:

As analgesic and anti-inflammatory in acute and chronic, painful and inflammatory processes of soft tissues and skeletal muscle. As analgesic and anti-inflammatory for bursitis, tendinitis, synovitis, tenosynovitis and lumbago. For analgesic action after dislocation, sprain, orthopedic manipulations, surgery and dental extraction. For uterine relaxation and post delivery analgesic for the mother who does not breastfeed. As analgesic in dysmenorrhea and after IUD insertion. As anti-inflammatory and analgesic drug in rheumatoid arthritis, osteoarthritis (degenerative arthritis), ankylosing spondylitis and gout.

## **EUROGESIC FORTE**

**Analgesic / Antipyretic / Anti-Inflammatory NSAID / Decongestant**

### **Composition:**

Each tablet contains:  
Naproxen sodium 550 mg  
Excipients q.s.

### **Presentations:**

10 tablet pack

### **Indications:**

As analgesic and anti-inflammatory in acute and chronic, painful and inflammatory processes of soft tissues and skeletal muscle. As analgesic and anti-inflammatory for bursitis, tendinitis, synovitis, tenosynovitis and lumbago. For analgesic action after dislocation, sprain, orthopedic manipulations, surgery and dental extraction. For uterine relaxation and post delivery analgesic for the mother who does not breastfeed. As analgesic in dysmenorrhea and after IUD insertion. As anti-inflammatory and analgesic drug in rheumatoid arthritis, osteoarthritis (degenerative arthritis), ankylosing spondylitis and gout.

# EUROGREL

## Antithrombotic

### Composition:

Each coated tablet contains:  
Clopidogrel (as bisulfate) 75 mg  
Excipients q.s.

### Presentations:

Hermetically sealed double aluminum blister pack with 35 tablets

### Indications:

EUROGREL is used to prevent vascular events (acute myocardial infarction, ischemic stroke) in patients with a history of symptomatic atherosclerosis. Also used to reduce atherosclerotic events in: myocardial infarction, acute coronary syndrome, ischemic stroke, peripheral arterial disease. Acute coronary syndrome (unstable angina/ non-Q-wave MI).

# EUROMICINA

## Antibiotic

### Composition:

Each tablet contains:  
Clarithromycin 250 mg  
Excipients q.s.

Clarithromycin 500 mg  
Excipients q.s.

Each 5 ml of reconstituted suspension contains:  
Clarithromycin 125 mg  
Excipients q.s.

Clarithromycin 250 mg  
Excipients q.s.

### Presentations:

Packs with 14 and 20 tablets of 250 or 500 mg.  
80 ml bottle of reconstituted suspension with 125 or 250 mg/5ml EUROMICINA, including solvent for reconstitution.

### Indications:

Clarithromycin is indicated for the treatment of infections caused by sensitive organisms. These infections include:

- Infection of the lower respiratory tract
- Infection of the upper respiratory tract.
- Skin and subcutaneous tissue infections.
- Infections caused or spread by *Mycobacterium avium* or intracellulare.
- Localized infections due to *Mycobacterium fortuitum* or *mycobacterium kanassi*.

Indicated together with Amoxicillin and omeprazol for the treatment of eradication of *Helicobacter pylori*, resulting in a reduction of duodenal ulcer recurrence.

# EUROMUCIL

## Laxative

### Composition:

Each tablespoon (11 g) contains:  
Psyllium hydrophilic mucilloid 3,4 g  
(Plantago ovata)  
Excipients q.s.

### Presentations:

Packs with 400 g of powder

### Indications:

EUROMUCIL is indicated in the treatment of constipation and bowel movement problems related to functional colonic disease or the chronic use of irritating laxatives. It is also used in dietary treatment for diverticulitis and hemorrhoids, or anal fissure. Moreover, it is used in regular or atonic constipation, constipation due to prolonged bed rest, and to regulate the evacuation in patients with iliac anus. At the beginning of treatment, the recommended starting dose is once daily, then gradually increased up to three times daily.

# EUROVIR

## Antiviral

### Composition:

Each tablet contains:

Acyclovir 400 mg

Excipients q.s.

Acyclovir 800 mg

Excipients q.s.

Each 5 ml of oral suspension contains:

Acyclovir 200 mg

Excipients q.s.

Each 5 ml of EUROVIR FORTE oral suspension contains:

Acyclovir 400 mg

Excipients q.s.

### Presentations:

EUROVIR 400 (Packs with 3 and 15 tablets of 400 mg)

EUROVIR 800 (Packs with 5 and 35 tablets of 800 mg)

EUROVIR Suspension (100 ml Bottle with 200 mg/5ml)

EUROVIR Suspension Forte: 100 ml bottle with 400 mg/5 ml

### Indications:

EUROVIR 400 is used in the treatment of herpes labialis and genital herpes, and in some forms of prophylaxis in immunocompromised patients.

EUROVIR 800 is used in: herpes zoster and suppression of recurrent genital herpes.

EUROVIR Suspension is specially used in children with: zoster-varicella, herpetic gingivostomatitis, severe herpes labialis, prophylaxis for herpes simplex infections in immunocompromised patients, and for suppression of recurrent herpes simplex infections in immunocompetent patients.

EUROVIR suspension Forte is used in the treatment of viral infections caused by herpes simplex virus types 1 and 2, herpes zoster and varicella-herpes zoster in immunodepressed patients.

## **EUROVIR CREMA**

### **Antiviral**

#### **Composition:**

Each gram of cream contains:

Acyclovir 50 mg

Excipients q.s.

#### **Presentations:**

EUROVIR Cream (5 and 15 gr. tubes)

#### **Indications:**

EUROVIR cream is used in: herpes labialis. It is also used as an oral coadjuvant therapy for other cutaneous herpetic infections



## **EVOCAZ CD**

### **Cholinesterase Inhibitor**

#### **Composition:**

Each dispersible tablet contains:

Donepezil Hydrochloride 5 mg  
Excipients q.s

Donepezil Hydrochloride 10 mg  
Excipients q.s.

#### **Presentations:**

EVOCAZ CD 5 mg: 30 dispersible tablets pack  
EVOCAZ CD 10 mg: 30 dispersible tablets pack

#### **Indications:**

EVOCAZ CD is indicated for the adjuvant therapy of Alzheimer-type dementia, mild to moderate nature.

## **FIBROLOW LIDOSE**

### **Antihyperlipidemic**

#### **Composition:**

Each capsule contains:  
Fenofibrate 200 mg  
Excipients q.s.

#### **Presentations:**

30 lidose capsule pack

#### **Indications:**

FIBROLOW Lidose is indicated as a complement to diet and non-pharmacological management (as exercise, weight reduction) in the next cases:

- Treatment of severe hypertriglyceridemia with or without low levels of HDL cholesterol.
- Mixed hyperlipidemia when statins are contraindicated or not tolerated by the patient.
- Mixed hyperlipidemia together with statin in patients with high cardiovascular risk, in whom triglycerides and HDL cholesterol have not been adequately controlled.

## **FIBROTINA LIDOSE**

### **Antihyperlipidemic**

#### **Composition:**

Each capsule contains:  
Fenofibrate 160 mg  
Pravastatin sodium 40 mg  
Excipients q.s.

#### **Presentations:**

30 lidose capsule pack

#### **Indications:**

FIBROTINA LIDOSE is indicated as a supplement for diets and other non-pharmacological treatments (eg, exercise, weight reduction) for the treatment of mixed hyperlipidemia in adult patients with a high cardiovascular risk, to reduce triglycerides and increase the level of HDL cholesterol when LDL cholesterol values are sufficiently controlled when treated with pravastatin 40 mg as monotherapy.

## **FIBROX XR**

### **Skeletal Muscle Relaxant**

#### **Composition:**

Each controlled release coated tablet contains:  
Cyclobenzaprine hydrochloride 10 mg  
Excipients q.s.

Cyclobenzaprine hydrochloride 20 mg  
Excipients q.s.

#### **Presentations:**

10 and 20 tablet pack

#### **Indications:**

Indicated for symptomatic treatment and as a coadjuvant treatment in rest and physical therapy for the relief of muscle spasms associated with acute musculoskeletal contractures.

# FINEX

## Antifungal

### Composition:

Each tablet contains:

Terbinafine (as hydrochloride) 250 mg

Excipients q.s.

Each 100 g of cream contains:

Terbinafine hydrochloride 1 g (1%)

Excipients q.s.

### Presentations:

FINEX tablets: packs with 14 and 28 tablets of 250 mg

FINEX cream: 15 g tube

### Indications:

FINEX tablets are indicated in fungal infections, where oral therapy is considered appropriate owing to the site, severity or extent of the infection.

FINEX is effective in the treatment of skin, hair and nails infections caused by dermatophytes such as Trichophyton (e.g., *T. rubrum*, *T. mentagrophytes*, *T. verrucosum*, *T. tonsurans*, *T. violaceum*), *Microsporum canis* and *Epidermophyton floccosum*.

Oral FINEX is indicated in the treatment of: tinea corporis, tinea cruris and tinea pedis, and cutaneous infections caused by yeast of the genus *Candida* (for example, *Candida albicans*). Onychomycosis caused by dermatophyte fungi. Oral terbinafine is not effective in pityriasis versicolor and vaginal candidiasis.

Topically applied terbinafine has a rapid onset of action and can be effective with a short duration of treatment. Less than 5% of the dose is absorbed after topical application to humans.

## **FOSVAL**

### **Antiresortive**

#### **Composition:**

Each tablet contains:  
Alendronic acid (as alendronato sodium trihydrate) 70 mg  
Excipients q.s.

#### **Presentations:**

FOSVAL 70 mg: 12 tablet blister calendar packs

#### **Indications:**

FOSVAL is indicated in the treatment and prevention of Osteoporosis in postmenopausal women.  
For the prevention of fractures, including hip and spine.  
Indicated for men and women in the treatment and prevention of glucocorticoid-induced osteoporosis.

## **FUNGIUM CREMA**

### **Antifungal**

#### **Composition:**

Each 100 g of cream contains:  
Ketoconazole 2 g (2%)  
Excipients q.s.

#### **Presentations:**

Tube with 15 g of dermal cream

#### **Indications:**

Topical treatment of skin infections: tinea corporis, tinea cruris, tinea manus and tinea pedis caused by *Trichophyton* sp., *Mocrosporium* sp., and *Epidermophyton* sp. Topical treatment of seborrheic dermatitis associated to infections caused by *Pityrosporum ovale*. Topical treatment of cutaneous candidiasis caused by *Candida albicans* and *Tinea versicolor* caused by *malassezia furfur*.

## **GESIX**

### **Nonsteroidal anti-inflammatory**

#### **Composition:**

Each capsule contains:  
Celecoxib 200 mg  
Excipients q.s.

#### **Presentations:**

10 and 30 capsule pack

#### **Indications:**

Gesix is indicated for:  
Symptomatic treatment of osteoarthritis and rheumatoid arthritis  
Relief of the signs and symptoms of ankylosing spondylitis  
Acute pain management  
Treatment of primary dysmenorrhea



# GINTOL

## Nutritional Supplement

### Composition:

Each capsule for oral administration contains:

Información nutricional	Por 100 g	1 porción	1 porción % *
Vitamina C (mg)	4168	40,00	67
Vitamina E (mg)	1250	12,00	60
Vitamina B1 (mg)	153	1,470	105
Vitamina B2 (mg)	178	1,710	107
Niacina (mg)	1667	16,00	89
Ácido pantonénico (mg)	625	6,00	60
Vitamina B6 (mg)	146	1,40	70
Biotina (mcg)	7918	76	25
Ácido fólico (mcg)	41675	400	200
Vitamina B12 (mcg)	260	2,5	250
Cobre (mg)	104	1,00	50
Hierro (mg)	1459	14,00	100
Zinc (mg)	1042	10,00	67
Manganeso (mg)	104	1,00	50
Selenio (mcg)	5730	55	79
Yodo (mcg)	15628	150	100
Cromo (mcg)	2084	20	57

\* Percentage according to recommended daily dose

### Presentations:

Pack with 30 soft capsules

## HEMOVAL

### Iron supplement

#### Composition:

Each chewable tablet contains:

Iron (as Iron III hydroxide polymallose complex) 100 mg

Excipients q.s.

Each ml of drop solution contains:

Iron (as Iron III hydroxide polymallose complex) 50 mg

Excipients q.s.

#### Presentations:

Chewable tablets: package of 42 tablets

Oral drops: 30 ml Bottle

#### Indications:

Prophylaxis and treatment of patients with iron-deficiency due to iron low intake, or poor absorption.

Uses

Iron deficiency states, such as anemia, pregnancy, and different growth stages.

## **HIALOF HIDRA**

### **Eye lubricant SP**

#### **Composition:**

Hialof Hidra does not contain preservatives.

Each 1 mL (20 drops) of sterile ophthalmic solution contains:

Sodium hyaluronate: 2 mg

Excipients q.s.

Each 100 mL of sterile ophthalmic solution contains:

Sodium hyaluronate: 0,2 g

Excipients q.s.

#### **Presentations:**

Sealed drop bottle with 10 mL ophthalmic solution

#### **Indications:**

It is indicated as an ocular lubricant, for the temporary relief of burning, foreign body sensation, stinging and dryness due to dry keratoconjunctivitis (external inflammatory conditions of the eye).

# HIDRIUM

**Diuretic**

## Composition:

Each tablet contains:  
Furosemide 40 mg  
Amiloride hydrochloride 5 mg  
Excipients q.s.

## Presentations:

20 scored tablet pack

## Indications:

HIDRIUM is indicated for the treatment of arterial hypertension, congestive heart failure, edematous states associated with heart failure, heart failure, nephrosis, corticosteroid or estrogen therapies, and cirrhosis-related ascites.

# IDON

## Antiemetic - Prokinetic

### Composition:

Each capsule contains:

Domperidone 10 mg

Excipients q.s.

Each 5 ml of oral suspension contains:

Domperidone 5 mg

Excipients q.s.

Each ml of drop solution contains:

Domperidone 10 mg

Excipients q.s.

### Presentations:

30 capsule pack

60 capsule pack

Bottle with 15 ml drops

Bottle with 100 ml of pediatric suspension

### Indications:

Adults:

- Relief of symptoms such as nausea and vomiting. Stomach burn sensation.

Children:

- Relief of symptoms of nausea and vomiting

## **IPSON / IPSON FORTE**

**Analgesic / Antipyretic**

### **Composition:**

Each 5ml of IPSON oral suspension contains:

Ibuprofen 100 mg

Excipients q.s.

Each 5 ml of IPSON FORTE oral suspension contains:

Ibuprofen 200 mg

Excipients q.s.

### **Presentations:**

IPSON: PET bottle with 120 ml of oral suspension

IPSON FORTE: PET bottle with 120 ml of oral suspension

### **Indications:**

IPSON is indicated for the treatment of inflammatory and painful processes of soft tissue and for febrile syndrome in patients over 6 months of age. Juvenile rheumatoid arthritis.

## **IPSON-D / IPSON-D FORTE**

**Analgesic / Antipyretic / Anti-Inflammatory NSAID**

### **Composition:**

Each 5 ml of IPSON D oral suspension contains:

Ibuprofen 100 mg

Pseudoephedrine hydrochloride 15 mg

Excipients q.s.

Each 5 ml of IPSON D FORTE oral suspension contains:

Ibuprofen 200 mg

Pseudoephedrine hydrochloride 30 mg

Excipients q.s.

Each IPSON D FORTE tablet contains:

Ibuprofen 400 mg

Pseudoephedrine hydrochloride 60 mg

Excipients q.s.

### **Presentations:**

Oral suspension: Bottle with 120 ml

Tablets: 10 tablet pack.

### **Indications:**

Ipson-D and Ipson-D Forte are indicated for transient relief of nasal congestion, headache and fever associated with flu or common cold. They are also useful in nasal congestion caused by infection or inflammation as well as in cases of sinusitis.

## **ITAN**

**Antiemetic - Prokinetic**

### **Composition:**

Each tablet contains:  
Metoclopramide (as hydrochloride) 10 mg  
Excipients q.s.

### **Presentations:**

30 tablet pack

### **Indications:**

Symptomatic treatment of postprandial pyrosis and gastric emptying delay in adults with gastroesophageal reflux. It is also used in the treatment of diabetic gastroparesia.

Uses:

Dysphagia; pyrosis; gastroparesia; gastric emptying delay; nausea and vomiting; regurgitation; gastroesophageal reflux; drug intolerance.



## **KEVAL**

### **Antimigraine**

#### **Composition:**

Each coated tablet contains:  
Eletriptan (as hydrobromide) 40 mg  
Excipients q.s.

#### **Presentations:**

Package with 2 or 6 coated tablets

#### **Indications:**

KEVAL® is indicated for the acute treatment of headache phase of migraine attacks with or without aura, for adults.

## **LATOF**

### **Antiglaucoma**

#### **Composition:**

Each 100 ml of ophthalmic solution contains:

Latanoprost 0,005 g

Excipients q.s.

#### **Presentations:**

Sealed drop bottle with 2.5 ml of sterile ophtalmic solution, with a controlled drop dispenser which aids in releasing drops making application easy and preventing drop loss and ensuring dose uniformity.

#### **Indications:**

Reduction of elevated intraocular pressure in patients with open-angle glaucoma and ocular hypertension.

## **LATOF SP**

### **Antiglaucoma**

#### **Composition:**

Latof SP does not contain preservatives.

Each 1 mL (32 drops) of ophthalmic solution of LATOF SP contains:

Latanoprost 0,05 mg

Excipients q.s.

Each 100 ml of ophthalmic solution of LATOF SP contains:

Latanoprost 0,005 g

Excipients q.s.

#### **Presentations:**

Sealed drop bottle with 2.5 ml of sterile ophthalmic solution

#### **Indications:**

Reduction of elevated intraocular pressure in patients with open angle glaucoma and ocular hypertension in patients with intolerance to other drugs that decrease intraocular pressure.

Reduction of elevated intraocular pressure in pediatric patients with elevated intraocular pressure and pediatric glaucoma.

## **LATOF-T**

### **Antiglaucoma**

#### **Composition:**

Each 100 ml of ophthalmic solution contains:

Latanoprost 0,005 g

Timolol maleate 0,5 g

Excipients q.s.

#### **Presentations:**

Sealed drop bottle with 2.5 ml of sterile ophtalmic solution, with a controlled drop dispenser which aids in releasing drops making application easy and preventing drop loss and ensuring dose uniformity

#### **Indications:**

Reduction of elevated intraocular pressure in patients with open-angle glaucoma and ocular hypertension.

## **LATOF-T SP**

### **Antiglaucoma**

#### **Composition:**

LATOF T SP does not contain preservatives.

Each 1 mL (33 drops) of ophthalmic solution of LATOF T SP contains:

Latanoprost 0,05 mg

Timolol: 5,00 mg

Excipients q.s.

Each 100 ml of ophthalmic solution of LATOF T SP contains:

Latanoprost 0,005 g

Timolol: 0,500 g

Excipients q.s.

#### **Presentations:**

Sealed drop bottle with 2.5 ml of sterile ophthalmic solution

#### **Indications:**

It is indicated for the reduction of elevated intraocular pressure (IOP) in patients with open angle glaucoma or ocular hypertension who respond insufficiently to topical agents that reduce IOP.

# LEOVAL

## Antiepileptic

### Composition:

Each coated tablet of LEOVAL 500 mg contains:  
Levetiracetam 500 mg  
Excipients q.s.

Each coated tablet of LEOVAL 1000 mg contains:  
Levetiracetam 1000 mg  
Excipients q.s.

### Presentations:

LEOVAL 500 mg: package with 30 coated tablets  
LEOVAL 1000 mg: package with 30 coated tablets

### Indications:

LEOVAL is indicated as monotherapy in the treatment of partial onset seizures, with or without secondary generalization in adults and adolescents over 16 years of age with a new diagnosis of epilepsy.

LEOVAL is indicated as concomitant therapy:

- In the treatment of partial onset seizures with or without secondary generalization in adults, adolescents and children over 4 years of age with epilepsy.
- In the treatment of myoclonic seizures in adults and adolescents over 12 years of age with Juvenile Myoclonic Epilepsy.
- In the treatment of primary generalized tonic-clonic seizures in adults and adolescents over 12 years of age with Generalized Idiopathic Epilepsy.

# LEOVAL INYECTABLE

## Antiepileptic

### Composition:

Each vial of 5 mL of solution contains:

Levetiracetam 500 mg

Excipients q.s.

### Presentations:

Pack containing 10 vials of 5 mL

### Indications:

LEOVAL is indicated as monotherapy in the treatment of partial onset seizures, with or without secondary generalization in adults and adolescents over 16 years of age with a new diagnosis of epilepsy.

LEOVAL is indicated as concomitant therapy:

- In the treatment of partial onset seizures with or without secondary generalization in adults, adolescents and children over 4 years of age with epilepsy.
- In the treatment of myoclonic seizures in adults and adolescents over 12 years of age with Juvenile Myoclonic Epilepsy.
- In the treatment of primary generalized tonic-clonic seizures in adults and adolescents over 12 years of age with Generalized Idiopathic Epilepsy.

Levetiracetam concentrate is an alternative for patients in whom oral administration is temporarily not feasible.

# LEOVAL SOLUCION

## Antiepileptic

### Composition:

Each mL of oral solution contains:

Levetiracetam 100 mg

Excipients q.s.

### Presentations:

Bottle with 300 mL of oral solution

### Indications:

LEOVAL is indicated as monotherapy in the treatment of partial onset seizures, with or without secondary generalization in adults and adolescents over 16 years of age with a new diagnosis of epilepsy.

LEOVAL is indicated as concomitant therapy:

- In the treatment of partial onset seizures with or without secondary generalization in adults, adolescents and children over 4 years of age with epilepsy.
- In the treatment of myoclonic seizures in adults and adolescents over 12 years of age with Juvenile Myoclonic Epilepsy.
- In the treatment of primary generalized tonic-clonic seizures in adults and adolescents over 12 years of age with Generalized Idiopathic Epilepsy.



## **LIFTER**

### **Erectile Dysfunction Agent**

#### **Composition:**

Each tablet contains:

Sildenafil (as citrate) 50 mg

Excipients q.s.

Sildenafil (as citrate) 100 mg

Excipients q.s.

#### **Presentations:**

50 mg tablets: packs with 1, 5 and 10 tablets

100 mg tablets: pack with 5 tablets

#### **Indications:**

For the treatment of erectile dysfunction.

## **LOMEX IV**

### **Antiulcer**

#### **Composition:**

Each vial of lyophilized contains:  
Omeprazole (as sodium salt) 40 mg  
Excipients q.s.

Each ampoule contains 10 ml of reconstitution solvent.

#### **Presentations:**

LOMEX 40 mg I.V. for injection:  
Combined pack containing one vial with omeprazole and one ampoule with 10 ml of solvent.

#### **Indications:**

LOMEX I.V.: prophylaxis of acid aspiration, prophylaxis of digestive hemorrhage in critical patients, upper gastrointestinal hemorrhage, Zollinger-Ellison syndrome and ulcer-related gastric retention.

## **LOWDEN**

### **Antihyperlipidemic**

#### **Composition:**

Each tablet contains:

Atorvastatin (as calcium salt) 10 mg

Excipients q.s.

Atorvastatin (as calcium salt) 20 mg

Excipients q.s.

Atorvastatin (as calcium salt) 40 mg

Excipients q.s.

#### **Presentations:**

LOWDEN 10 mg x 30 and 60 scored tablets

LOWDEN 20 mg x 30 scored tablets

LOWDEN 40 mg x 30 scored tablets

All tablets are packed in double laminated blister packs to protect the active substance

#### **Indications:**

LOWDEN is indicated for the reduction of total cholesterol and LDL cholesterol in patients with hypercholesterolemia as adjunctive therapy to diet.

## **LUDIUM 20 MG**

### **Phosphodiesterase Type 5 Inhibitor**

#### **Composition:**

Each coated tablet contains:

Tadalafil 20 mg

Excipients q.s.

#### **Presentations:**

Package with 1 and 4 coated tablets

#### **Indications:**

LUDIUM 20 mg is indicated in:

- The treatment of erectile dysfunction in adult men. In order for LUDIUM to be effective, sexual stimulation is necessary.

The use of LUDIUM is not indicated in women.

## **LUDIUM 5 MG**

**Phosphodiesterase Type 5 Inhibitor**

### **Composition:**

Each coated tablet contains:

Tadalafil 5 mg

Excipients q.s.

### **Presentations:**

Package with 30 coated tablets

### **Indications:**

LUDIUM 5 mg is indicated in:

- Treatment of erectile dysfunction in adult men. In order for LUDIUM to be effective, sexual stimulation is necessary.
- Treatment of the signs and symptoms of benign prostatic hyperplasia in adult men.

The use of LUDIUM is not indicated in women.

# LUKANEX

## Antiasthmatic

### Composition:

Each chewable tablet contains:

Montelukast sodium: 4 mg

Excipients q.s.

Montelukast sodium: 5 mg

Excipients q.s.

Each coated tablet contains:

Montelukast: 10 mg

Excipients q.s.

Each sachet contains:

Montelukast: 4 mg

Excipients q.s.

### Presentations:

4 mg granulated Lukanex: 40 sachets pack

4 mg Lukanex chewable tablets: 40 chewable tablet pack

5 mg Lukanex chewable tablets: 40 chewable tablet pack

10 mg Lukanex coated tablets: 40 coated tablets pack

### Indications:

Montelukast is indicated in:

- Prophylaxis and chronic treatment of asthma in adults and pediatric patients from 12 months of age
- Prevention of exercise-induced bronchoconstriction in adult and adolescent patients older than 15 years of age
- Relief of symptoms of seasonal allergic rhinitis in adult and adolescent patients from 2 years of age
- Relief of symptoms of perennial allergic rhinitis in adult patients and children from 6 months of age

## **MENTANIA**

**Vaso-dilatant / Brain Stimulant**

### **Composition:**

Each soft capsule contains:

Standardized dry extract of Gingko biloba leaves 60 mg

(Equivalent to 14,4 mg of flavonglycosides)

Standardized dry extract of Panax ginseng root C.A.Meyer 100 mg

(Equivalent to 4 mg of glycosides)

Excipients q.s.

### **Presentations:**

30 soft capsule packs

### **Indications:**

Treatment of cerebral vascular insufficiency and some its manifestations such as cognitive deficit, mood swings and the like.

# MIGRAX

## Antijaquecoso

### Composition:

Each coated tablet contains:

Acetaminophen 450 mg

Caffeine 40 mg

Dihydroergotamine mesylate 1 mg

Excipients q.s.

### Presentations:

10 tablet packs

### Indications:

MIGRAX is indicated for the symptomatic treatment of vascular headaches.



## **MIGTAL**

### **Antimigraine**

#### **Composition:**

Each coated tablet contains:  
Naratriptan (as hydrochloride) 2,5 mg  
Excipients q.s.

#### **Presentations:**

2, 6 and 12 tablet Pack

#### **Indications:**

For the treatment of headache – migraine: Indicated for the treatment of acute migraine attacks with or without aura in adults.

# MOXAVAL

## Antibiotic

### Composition:

Each coated tablet contains:  
Moxifloxacin (as hydrochloride) 400 mg  
Excipients q.s.

### Presentations:

7 and 10 coated tablet packs

### Indications:

Moxifloxacin is indicated for the treatment of adults over 18 years of age with upper and lower respiratory tract infections such as acute sinusitis; acute exacerbation of chronic bronchitis; extra hospital pneumonia. In addition, it is indicated for cutaneous infections and soft tissue infections.

## **MOXOF**

### **Ophthalmic Antibiotic**

#### **Composition:**

Each 100 ml of ophthalmic solution contains:  
Moxifloxacin (as hydrochloride) 0,5 g  
Excipients q.s.

#### **Presentations:**

Sealed drop-dispenser bottle with 5 mL of sterile ophthalmic solution

#### **Indications:**

Moxof is indicated for the treatment of bacterial conjunctivitis caused by susceptible microorganism strains.

# MUXELIX

## Mucolytic

### Composition:

Each 100 ml of oral solution contains:  
Hedera helix L. dry extract (5-7,5 : 1) 0,7 g  
(Equivalent to no less than 70 mg hederacoside C)  
Excipients q.s.

### Presentations:

120 ml syrup bottle

### Indications:

Treatment of inflammatory bronchial disease symptoms associated with cough.

# MUXOL

## Mucolytic

### Composition:

Each tablet contains:  
Ambroxol hydrochloride 30 mg  
Excipients q.s.

Each 5 ml of adult oral solution contains:  
Ambroxol hydrochloride 30 mg  
Excipients q.s.

Each 5 ml of pediatric oral solution contains:  
Ambroxol hydrochloride 15 mg  
Excipients q.s.

### Presentations:

20 tablet pack  
100 ml adult syrup bottle  
100 ml pediatric syrup bottle

### Indications:

Ambroxol is indicated as mucolytic expectorant agent for the relief of productive cough caused by flu states.

## **NEUROVAL CD**

### **Antianxiety**

#### **Composition:**

Each dispersible tablet contains:

Clotiazepam 5 mg

Excipients q.s.

Each dispersible tablet contains:

Clotiazepam 10 mg

Excipients q.s.

#### **Presentations:**

NEUROVAL®CD 5 mg: 30 dispersible tablet packs

NEUROVAL®CD 10 mg: 30 dispersible tablet packs

#### **Indications:**

Treatment of generalized anxiety

## NEURUM

### Antiepileptic

#### Composition:

Each capsule contains:  
Pregabalin 50 mg  
Excipients q.s.

Pregabalin 75 mg  
Excipients q.s.

Pregabalin 150 mg  
Excipients q.s.

#### Presentations:

Neurum 50 mg x 30 capsule pack  
Neurum 75 mg x 30 y 40 capsule pack  
Neurum 150 mg x 30 capsule pack

#### Indications:

Treatment of neurophatic pain in adults. Adjuvant therapy for patients 12 years of age and older, with or without secondary generalization.

## **NICODROPS**

### **Ophthalmic Decongestant Drops**

#### **Composition:**

Each 100 ml of ophthalmic solution contains:

Naphazoline hydrochloride 0,025 g

Dextran-70 0,100 g

Hypromellose 0,300 g

Excipients q.s.

#### **Presentations:**

10 ml sterile ophthalmic solution bottle

#### **Indications:**

Transient relief of eye redness due to minor ocular irritation. It is indicated in all conditions in which relief of irritation, burning, pruritus and/or ocular congestion is required, for example: dust, smoke, air pollution, sun exposure, wind, contact lenses, allergy, cold or swimming.



## **NICOL**

### **Oral Contraceptive**

#### **Composition:**

Each active, white coated tablet contains:

Dienogest 2.00 mg

Ethinyl estradiol 0.03 mg

Excipients q.s.

Each placebo, green coated tablet contains:

Excipients q.s.

#### **Presentations:**

Pack containing 28 coated tablets

#### **Indications:**

Hormonal contraception.

Treatment of moderately severe acne, which is refractory to local treatment in women in whom the administration of oral contraceptives is not contraindicated.

The decision to prescribe Nicol should take into account the current risk factors of each individual woman, specifically those of venous thromboembolism and how the risk of venous thromboembolism compares with that of other combined hormonal contraceptives.

# NICOTEARS

## Ophthalmic Drops substitute

### Composition:

Each ml of sterile ophthalmic solution contains:

Hypromellose 3 mg

Dextran-70 1 mg

Excipients q.s.

### Presentations:

Sealed drop-dispenser bottle with 20 ml sterile ophthalmic solution

### Indications:

NICOTEARS is indicated as a lubricating and moistening agent for the eye in the symptomatic treatment of dry eye. It is also indicated in the irritation caused by contact lenses; in Keratitis caused by exposure to irritants; decreased corneal sensitivity; burning; photophobia; conjunctival hyperemia, and generally in recurrent corneal and/or conjunctival erosions and irritations.

## **NICOTEARS GEL**

**Substitute for tears**

### **Composition:**

Each g of sterile ophthalmic gel contains:

Carbomer 940 1,5 mg

Manitol 46 mg

Excipients q.s.

### **Presentations:**

Sealed tube with 5 g of sterile ophthalmic gel.

### **Indications:**

Relief of the dry eye due to environmental agents such as wind, cigarette smoke, pollution, and solar radiation, or additionally, by physiological conditions manifested by a deficit in tear production. It is also indicated in irritation caused by hard contact lenses; exposure Keratitis, burning, photophobia; conjunctival hyperemia and generally in superior corneal irritations.

## **NIRVAN**

### **Nonbenzodiazepine Hypnotic**

#### **Composition:**

Each coated tablet contains:

Eszopiclone 2 mg

Excipients q.s.

Eszopiclone 3 mg

Excipients q.s.

#### **Presentations:**

40 scored coated tablets pack

#### **Indications:**

Indicated for the treatment of temporary or chronic insomnia

# NOVOTEARS

## Ophthalmic Decongestant Drops

### Composition:

Each 100 ml of sterile ophthalmic solution contains:

Naphazoline hydrochloride 0,012 g

Hypromellose 0,3 g

Dextran-70 0,1 g

Excipients q.s.

### Presentations:

Sealed drop-dispenser bottle with 10 ml sterile ophthalmic solution

### Indications:

Transient relief of eye redness due to minor ocular irritations such as burning sensation, itching, and/or ocular congestion, due to dust, smoke, air pollution, sun exposure, wind, contact lenses, allergy or swimming, winds.

# **NYSKIN**

## **Antiacne**

### **Composition:**

Each 100 g of dermal gel 0,3% contains:

Adapalene 0,3 g

Excipients q.s.

Each 100 g of dermal gel 0,1% contains:

Adapalene 0,1 g

Excipients q.s.

Each 100 g of dermal cream 0,1% contains:

Adapalene 0,1 g

Excipients q.s.

### **Presentations:**

35 g cream or gel tubes

### **Indications:**

Adapalene is indicated for the topical treatment of acne vulgaris.

# OBEXOL

## Anorexigenic Agent

### Composition:

Each capsule contains:  
Phentermine Hydrochloride 18,75 or 37,5 mg  
Excipients q.s.

### Presentations:

30 capsule pack

### Indications:

OBEXOL is indicated for short-term treatment of exogenous obesity together with diet to reduce body weight, based on exercise, modification of eating habits and calorie restriction. The treatment is suitable for patients with a body mass index (BMI) greater than or equal to 30 kg / m<sup>2</sup> or greater than or equal to 27 kg / m<sup>2</sup> in the presence of risk factors such as hypertension, diabetes and hyperlipidemia.

# OFTABIÓTICO

## Ophthalmic Antibiotic

### Composition:

Each ml of sterile ophthalmic solution contains:

Polymyxin B (as sulfate) 5000 I.U.

Neomycin (as sulfate) 1,7 mg

Gramicidin 0,025 mg

Excipients q.s.

### Presentations:

Sealed drop bottle with 10 ml of sterile ophthalmic solution

### Indications:

OFTABIOTICO is indicated for short-term treatment of superficial external ocular infections caused by sensitive organisms.



# OFTABIÓTICO UNGÜENTO

## Ophthalmic Antibiotic

### Composition:

Each g of sterile ophthalmic ointment contains:

Polymyxin B (as sulfate) 6000 I.U.

Neomycin (as sulfate) 3,5 mg

Bacitracin 400 I.U.

Excipients q.s.

### Presentations:

3.5 g ointment tubes.

### Indications:

OFTABIOTICO ointment is indicated for short-term treatment of superficial external ocular infections caused by sensitive organisms.

## OFTAFILM

### Eye lubricant

#### Composition:

Each 100 ml of ophthalmic solution contains:

Sodium hyaluronate 0,400 g

Excipients q.s.

#### Presentations:

Sealed drop bottle with 10 ml of sterile ophthalmic solution

#### Indications:

It is indicated as ocular lubricant for the transient relief of burning sensation, foreign body sensation, itching and dryness due to keratoconjunctivitis sicca.

Uses: keratoconjunctivitis sicca, exposure keratitis, neuroparalitic keratitis, mild ocular irritation by sun rays, dust, air, chlorinated waters, weak chemical agents, weak chemical agents, and exposure to strong light.

## **OFTAFILM SP**

### **Eye lubricant SP**

#### **Composition:**

Each 100 ml of sterile ophthalmic solution contains:  
Sodium hyaluronate 0,400 g  
Excipients q.s.

#### **Presentations:**

Sealed drop bottle with 10 mL ophthalmic solution

#### **Indications:**

It is indicated as ocular lubricant for the transient relief of burning sensation, foreign body sensation, itching and dryness due to dry keratoconjunctivitis.

Pediatric use: Do not use, safety and efficacy of this product has not been established in pediatric patients.

Geriatric use: No significant differences were observed in clinical safety and efficacy in elderly patients compared to younger people.

# OFTAGEN

## Ophthalmic Antibiotic

### Composition:

Each ml of sterile ophthalmic solution contains:

Gentamicin (as sulfate) 3 mg

Excipients q.s.

Each g of ophthalmic ointment contains:

Gentamicin (as sulfate) 3 mg

Excipients q.s.

### Presentations:

Sealed drop bottle with 5 ml of sterile ophthalmic solution.

3.5g ointment tube.

### Indications:

OFTAGEN is indicated for the treatment of bacterial infections of the surface structure of the eye caused by microorganisms sensitive to gentamicin, for example, conjunctivitis, keratitis, keratoconjunctivitis, corneal ulcer, blepharitis, blepharoconjunctivitis, meibomianitis, and dacriocystitis.

It is also indicated in the prophylaxis of postoperative ocular infections.

# OFTAGEN COMPUESTO

## Ophthalmic Antibiotic / Corticosteroid

### Composition:

Each ml of sterile ophthalmic solution contains:

Gentamicin (as sulfate) 3 mg

Bethamethasone disodium fosfate 1 mg

Excipients q.s.

Each g of ophthalmic ointment contains:

Gentamicin (as sulfate) 3 mg

Bethamethasone disodium phosphate 1 mg

Excipients q.s.

### Presentations:

Sealed drop bottle with 5 ml of sterile ophthalmic solution

3.5 g ointment tube

### Indications:

OFTAGEN is indicated for the treatment of bacterial infections of the surface structure of the eye caused by microorganisms sensitive to gentamicin; associated with inflammatory and allergic conditions such as conjunctivitis, keratitis, and keratoconjunctivitis. Indicated as prophylaxis of postoperative ocular infections of the eye.

## **OFTALER / OFTALER FORTE**

### **Ophthalmic Antihistamine**

#### **Composition:**

Each 100 ml of ophthalmic solution contains:

Ketotifen (as fumarate) 0,025 g (0,025%)

Excipients q.s.

Ketotifen (as fumarate) 0,050 g (0,050%)

Excipients q.s.

#### **Presentations:**

Sealed drop bottle with 10 ml of sterile ophthalmic solution 0.05% and 0.025%.

#### **Indications:**

Relief of the signs and symptoms of allergic conjunctivitis.

## **OFTALIRIO**

**Ophthalmic Decongestant / Antihistamine**

### **Composition:**

Each 100 ml of sterile ophthalmic solution contains:  
Naphazoline hydrochloride 0,05 g  
Antazoline fosfate 0,50 g  
Excipients q.s.

### **Presentations:**

Sealed drop bottle with 10 ml of sterile ophthalmic solution.

### **Indications:**

The onset of action with OFTALIRIO can be noted within few minutes after topical application and its effect lasts for 2 -3 hours. Rapid relief of pruritus and ocular congestive symptoms associated with irritating, allergic and inflammatory conditions.

## OFTASONA-N

### Ophthalmic Antibiotic / Corticosteroid

#### Composition:

Each 100 ml of sterile ophthalmic solution contains:

Bethamethasone disodium phosphate 0,1 g

Neomycin (as sulfate) 0,35 g

Excipients q.s.

Each 100 g of ophthalmic ointment contains:

Bethamethasone disodium phosphate 0,1 g

Neomycin (as sulfate) 0,35 g

Excipients q.s.

#### Presentations:

Sealed drop bottle with 10 ml of sterile ophthalmic solution

3 g ointment tube.

#### Indications:

OFTASONA-N is indicated for the treatment of infectious processes caused by germs sensitive to neomycin accompanied by an inflammatory component.



## **OFTASONA-P**

### **Ophthalmic Corticosteroid**

#### **Composition:**

Each 100 ml of sterile ophthalmic solution contains:  
Bethamethasone disodium phosphate 0,1 g  
Excipients q.s.

#### **Presentations:**

Sealed drop bottle with 10 ml of sterile ophthalmic solution.

#### **Indications:**

Indicated for the treatment of allergic conjunctivitis, inflammatory conditions of the anterior segment of the eye, caused by thermal, chemical, postoperative, and radiation factors.

## **OFTAVIR**

### **Ophthalmic Antiviral**

#### **Composition:**

Each 100 g of ophthalmic ointment contains:

Acyclovir 3 g

Excipients q.s.

#### **Presentations:**

3.5 g tube with sterile ophthalmic ointment

#### **Indications:**

OFTAVIR is indicated in the selective treatment of Herpes simplex keratitis. Coadjuvant therapy in systemic treatment of Herpes Zoster with ocular involvement.

# OFTAVITA

## Polivitamin with minerals

### Composition:

Each coated tablet contains:

Vitamin A (As beta carotene)	3300 IU	(2,00 mg)
Vitamin C (As ascorbic acid)	200,00 mg	
Vitamin E (As DL- <sup>1</sup> Tocopheryl acetate)	75,00 IU	(75,00 mg)
Vitamin B2	5,00 mg	
Zinc (As acetate dihydrate)	30,00 mg	
Selenium (As amino acid complex)	0,02 mg	
Copper (As glycinate chelate)	2,00 mg	
Manganese (As glycinate chelate)	5,00 mg	
Lutein / zeaxanthin	2,00 mg	
Excipients q.s.		

### Presentations:

OFTAVITA®: 30 coated tablets pack

### Indications:

Vitamin and mineral deficiency states

## **OFTIC**

### **Ophthalmic Analgesic - Anti-Inflammatory NSAID**

#### **Composition:**

Each 100 ml of sterile ophthalmic solution contains:

Diclofenac sodium 0,1 g

Excipients q.s.

#### **Presentations:**

10 ml sealed pack with sterile ophthalmic solution.

#### **Indications:**

OFTIC is used in the treatment of inflammation in patients undergoing cataract surgery.

## OFTOL / OFTOL FORTE

### Ophthalmic Corticosteroid

#### Composition:

100 mL of OFTOL ophthalmic suspension contain:  
0.200 g Loteprednol Etabonate  
Excipients c.s.

100 mL of OFTOL FORTE ophthalmic suspension contain:  
0.500 g Loteprednol Etabonate  
Excipients c.s.

#### Presentations:

Sealed drop bottle containing 5 mL of sterile ophthalmic suspension.

#### Indications:

The physician will indicate the concentration and dose of the drug to be used according to the nature and intensity of the clinical condition. Generally, OFTOL FORTE is used in the treatment of inflammatory conditions of the palpebral and bulbar conjunctiva, cornea and anterior segment of the globe that respond to steroids such as allergic conjunctivitis, acne rosacea, superficial punctate keratitis, herpes zoster keratitis, iritis, cyclitis, selected infective conjunctivitis, when the potential damage of steroid use has been accepted, in order to obtain a decrease in edema and inflammation. It is also used in the treatment of inflammation after ocular surgery. OFTOL is usually indicated for transient relief of signs and symptoms of seasonal allergic conjunctivitis.

# OFTOL PLUS

## Ophthalmic Antibiotic / Corticosteroid

### Composition:

Each 100 ml of Ophthalmic Solution contains:  
Loteprednol Etabonate 0,5 g  
Tobramycin 0,3 g  
Excipients q.s.

### Presentations:

Sealed drop bottle with 5 mL of sterile ophthalmic solution

### Indications:

The association is indicated for the treatment of inflammatory eye conditions who present superficial bacterial infection or when there is a risk of bacterial ocular infection.

Corticosteroids are indicated for inflammation of the conjunctiva of eyelids and bulbar conjunctiva, cornea, and anterior segment of the eye including allergic conjunctivitis, acne rosacea, superficial punctate keratitis, herpes zoster keratitis, iritis, cyclitis.

It is also used in cases of infectious conjunctivitis when it is beneficial to reduce edema and inflammation and at the same time when the risk of using topical corticosteroids agents is justified.

It is used to treat chronic anterior uveitis and cornea damage due to chemicals components, radiation, burns, or penetration of foreign bodies.

The use of Loteprednol combined with tobramycin (antibiotic) is indicated where the risk of superficial ocular infection is high or where there is a potential risk of bacterial growth in the eye.

## OLOF

### Decongestant / Antihistamine

#### Composition:

Each 1 mL (28 drops) of ophthalmic solution contains:

Olopatadine (Hydrochloride) 2 mg

Excipients: q.s.

Each 100 mL of ophthalmic solution contains:

Olopatadine (Hydrochloride) 0,2 g

Excipients: q.s.

#### Presentations:

Sealed drop bottle containing 5 mL of sterile ophthalmic suspension.

#### Indications:

Indicated for the treatment of ocular itching associated with allergic conjunctivitis.

# OTICUM

**Otic Antibiotic / Corticosteroid**

## Composition:

Each 100 ml of otic solution contains:

Polymyxin B (as sulfate) 1000000 I.U.

Neomycin (as sulfate) 0,35 g

Bethamethasone disodium phosphate 0,1 g

Lidocaine hydrochloride 2 g

Excipients q.s.

## Presentations:

5 ml sterile solution pack

## Indications:

OTICUM is indicated for the treatment of external otitis accompanied by pain and inflammation, caused by sensitive bacteria. Complementary treatment of acute and chronic otitis media.

Indicated as a coadjuvant in the parenteral antibiotic treatment and other therapeutic measures, and in cavity protection following mastoidectomy and fenestration.



# **PAXON**

## **Antianxiety**

### **Composition:**

Each tablet contains:

Buspirone hydrochloride 5 mg

Excipients q.s.

Buspirone hydrochloride 10 mg

Excipients q.s.

### **Presentations:**

Packs with 20 scored tablets of 5 mg or 10 mg

### **Indications:**

PAXON is indicated for the treatment of acute and chronic anxiety disorders with the following symptoms: anxiety, unsettledness, internal restlessness, and tension states.

## **PERTIUM**

### **Antihypertensive**

#### **Composition:**

Each tablet contains:  
Nebivolol (as hydrochloride) 5 mg  
Excipients q.s.

#### **Presentations:**

Package of 42 tablets

#### **Indications:**

Treatment of essential hypertension alone or in combination with other antihypertensive agents. Treatment of stable, mild, moderate, and severe heart failure in addition to standard therapies (eg diuretics, ACE inhibitors and angiotensin II antagonists) in elderly patients > 70 years of age.

## **PILOCARPINA**

**Antiglaucoma / Direct Acting Miotic**

### **Composition:**

Each ml of sterile ophthalmic solution contains:

Pilocarpine hydrochloride 20 mg (2%)  
Excipients q.s.

Pilocarpine hydrochloride 40 mg (4%)  
Excipients q.s.

### **Presentations:**

Sealed drop bottle with 10 ml of sterile ophthalmic solution 2% or 4%.

### **Indications:**

Antagonist of the mydriatic and cycloplegic effects of atropine. Open angle glaucoma. Narrow angle glaucoma.

## **PROSTOP-D**

**Alpha adrenergic receptor antagonist**

### **Composition:**

Each capsule contains:

Dutasteride 0.5 mg

Tamsulosin (as hydrochloride) 0.4 mg

Excipients q.s.

### **Presentations:**

Pack with 30 capsules

### **Indications:**

For the treatment of the symptoms of benign prostatic hyperplasia (BPH) in men with an enlarged prostate.

## **PROTIUM - F**

### **Nutritional Supplement**

#### **Composition:**

Each capsule contains:

*Lactobacillus acidophilus* La-14® (4x10<sup>9</sup> UFC)

*Lactobacillus rhamnosus* HN001® (1x10<sup>9</sup> UFC)

Excipients q.s.

#### **Presentations:**

30 capsules pack

## **PROTIUM - I**

### **Nutritional Supplement**

#### **Composition:**

Each capsule contains:

*Lactobacillus acidophilus* NCFM® + *Bifidobacterium lactis* Bi-07® (1\*10<sup>10</sup> UFC)

Excipients q.s.

#### **Presentations:**

30 capsules pack

## **PROTIUM - T**

### **Nutritional Supplement**

#### **Composition:**

Each capsule contains:

*Bifidobacterium lactis* HN019® + *Lactobacillus acidophilus* NCFM® (1\*10<sup>10</sup> UFC)

Excipients q.s.

#### **Presentations:**

30 capsules pack

## **PRUVAL**

### **Other drugs for constipation**

#### **Composition:**

Each coated tablet 1 mg PRUVAL contains:  
Prucalopride (as succinate) 1 mg  
Excipients q.s.

Each coated tablet 2 mg PRUVAL contains:  
Prucalopride (as succinate) 2 mg  
Excipients q.s.

#### **Presentations:**

Pruval 1 mg: 30 tablet packs  
Pruval 2 mg: 30 tablet packs

#### **Indications:**

PRUVAL is indicated for the symptomatic treatment of chronic constipation in women for whom laxatives have not provided adequate relief.



# QUETIUM

## Antipsychotic

### Composition:

Each coated tablet of QUETIUM® 25 mg contains:  
Quetiapine (fumarate) 25 mg  
Excipients q.s.

Each coated tablet of QUETIUM® 100 mg contains:  
Quetiapine (fumarate) 100 mg  
Excipients q.s.

Each coated tablet of QUETIUM® 200 mg contains:  
Quetiapine (fumarate) 200 mg  
Excipients q.s.

Each coated tablet of QUETIUM® 300 mg contains:  
Quetiapine (fumarate) 300 mg  
Excipients q.s.

### Presentations:

QUETIUM® 25 mg: package of 30 coated tablets  
QUETIUM® 100 mg: package of 30 coated tablets  
QUETIUM® 200 mg: package of 30 coated tablets  
QUETIUM® 300 mg: package of 30 coated tablets

### Indications:

QUETIUM® is indicated in:

- Schizophrenia
- Manic episodes associated with bipolar disorder
- Depressive episodes associated with bipolar disorder
- Preventing recurrences in the treatment of maintenance of bipolar disorder (manic, mixed or depressive episodes) in combination with lithium or valproate

# REALTA

## Antidepressant

### Composition:

Each capsule with enteric-coated granules of Realta 30 mg contains:

Duloxetine: 30 mg

Excipients q.s.

Each capsule with enteric-coated granules of Realta 60 mg contains:

Duloxetine: 60 mg

Excipients q.s.

### Presentations:

Realta 30 mg: 30 capsule pack

Realta 60 mg: 30 capsule pack

### Indications:

Realta is indicated in:

- Major Depressive Disorder
- Generalized Anxiety Disorder
- Diabetic Peripheral Neuropathic Pain
- Fibromyalgia
- Musculoskeletal chronic pain

# REMSIMA

## Tumor necrosis factor inhibitor

### Composition:

One vial contains 100 mg of infliximab. After reconstitution each mL contains 10 mg of infliximab.

### Presentations:

1 vial of 100 mg

### Indications:

#### Rheumatoid arthritis:

Remsima, in combination with methotrexate, is indicated for the reduction of signs and symptoms as well as the improvement in physical function in:

- adult patients with active disease when the response to disease-modifying antirheumatic drugs (DMARDs), including methotrexate, has been inadequate.
- adult patients with severe, active and progressive disease not previously treated with methotrexate or other DMARDs.

In these patient populations, a reduction in the rate of the progression of joint damage, as measured by X-ray, has been demonstrated.

#### Adult Crohn's disease:

Remsima is indicated for:

- treatment of moderately to severely active Crohn's disease, in adult patients who have not responded despite a full and adequate course of therapy with a corticosteroid and/or an immunosuppressant; or who are intolerant to or have medical contraindications for such therapies.
- treatment of fistulising, active Crohn's disease, in adult patients who have not responded despite a full and adequate course of therapy with conventional treatment (including antibiotics, drainage and immunosuppressive therapy).

#### Paediatric Crohn's disease:

Remsima is indicated for treatment of severe, active Crohn's disease in children and adolescents aged 6 to 17 years, who have not responded to conventional therapy including a corticosteroid, an immunomodulator and primary nutrition therapy; or who are intolerant to or have contraindications for such therapies. Infliximab has been studied only in combination with conventional immunosuppressive therapy.

#### Ulcerative colitis:

Remsima is indicated for treatment of moderately to severely active ulcerative colitis in adult patients who have had an inadequate response to conventional therapy including corticosteroids and 6-mercaptopurine (6-MP) or azathioprine (AZA), or who are intolerant to or have medical contraindications for such therapies.

#### Ankylosing spondylitis:

Remsima is indicated for treatment of severe, active ankylosing spondylitis, in adult patients who have responded inadequately to conventional therapy.

#### Psoriatic arthritis:

Remsima is indicated for treatment of active and progressive psoriatic arthritis in adult patients when the response to previous DMARD therapy has been inadequate.

Remsima should be administered:

- in combination with methotrexate
- or alone in patients who show intolerance to methotrexate or for whom methotrexate is contraindicated.

Infliximab has been shown to improve physical function in patients with psoriatic arthritis, and to reduce the rate of progression of peripheral joint damage as measured by X-ray in patients with polyarticular symmetrical subtypes of the disease.

**Psoriasis:**

Remsima is indicated for treatment of moderate to severe plaque psoriasis in adult patients who failed to respond to, or who have a contraindication to, or are intolerant to other systemic therapy including cyclosporine, methotrexate or psoralen ultra-violet A (PUVA).

## **RINOVAL**

### **Nasal corticosteroid**

#### **Composition:**

Each dose contains:

Mometasone furoate (as monohydrate) 50 mcg

Excipients q.s.

#### **Presentations:**

Nasal spray bottle with 120 doses of mometasone furoate 50 mcg/dose.

#### **Indications:**

- Treatment of the symptoms of seasonal and perennial rhinitis in adults and pediatric patients from 2 years of age.
- Prophylaxis of the nasal symptoms of allergic rhinitis in adults and children older than 12 years of age, 2 to 4 weeks prior to the commencement of the exposure to allergens.
- Indicated in adults and patients over 12 years of age as adjuvant to the antibiotic therapy in the treatment of acute sinusitis episodes.
- Treatment of nasal polyps in patients from 18 years of age.

## **RIVOXAXAN 10 mg**

### **Platelet Aggregation Inhibitor**

#### **Composition:**

Each Rivoxa film-coated tablet contains:

Rivaroxaban 10 mg

Excipients: q.s.

#### **Presentations:**

Rivoxa 10 mg: Envase con 10 comprimidos recubiertos

Bioequivalente: este producto ha demostrado su equivalencia terapéutica.

#### **Indications:**

Prevention of venous thromboembolism (VTE) in patients undergoing major orthopedic surgery of the lower extremities.

Treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE), and prevention of recurrences of DVT and PE in adults.

## **RIVOXÁ 15 mg y 20 mg**

### **Platelet Aggregation Inhibitor**

#### **Composition:**

Each Rivoxa film-coated tablet contains:

Rivaroxaban 15 or 20 mg

Excipients: q.s.

#### **Presentations:**

Rivoxa 15 mg: Package with 30 coated tablets

Rivoxa 20 mg: Package with 30 coated tablets

Bioequivalent: This product has demonstrated its therapeutic equivalence

#### **Indications:**

Prevention of cerebrovascular accident and systemic embolism in adult patients with nonvalvular atrial fibrillation, with one or more risk factors, such as congestive heart failure, hypertension, age  $\geq$  75 years, diabetes mellitus, previous cerebrovascular accident or transient ischemic attack.

Treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE), and prevention of recurrences of DVT and PE in adults.

# RUX

## Antihyperlipidemic

### Composition:

Each film coated tablet contains:

Rosuvastatin (as calcium salt) 5 mg  
Excipients q.s.

Rosuvastatin (as calcium salt) 10 mg  
Excipients q.s.

Rosuvastatin (as calcium salt) 20 mg  
Excipients q.s.

### Presentations:

RUX 5 mg: 30 coated tablets pack  
RUX 10 mg: 30 and 60 coated tablets packs  
RUX 20 mg: 30 coated tablets pack

### Indications:

Rosuvastatin is indicated for the treatment of:

Adults:

#### 1. Primary hyperlipidemia and mixed dyslipidemia

Rosuvastatin is indicated as adjunctive therapy to diet to reduce elevated total cholesterol (TC), LDL-C, ApoB, non-HDL cholesterol and triglycerides and to increase HDL-C in adult patients with primary hyperlipidemia or mixed dyslipidemia.

#### 2. Hypertriglyceridemia

Rosuvastatin is indicated as adjunctive therapy to diet for the treatment of patients with hypertriglyceridemia.

#### 3. Type III familial hyperlipoproteinemia.

Rosuvastatin is indicated as adjunctive therapy to diet for the treatment of patients with Type III familial hyperlipoproteinemia (a genetic disease characterized by hypertriglyceridemia, hypercholesterolemia, and the presence of cholesterol-rich, very low density lipoproteins in plasma).

#### 4. Homozygous familial hypercholesterolemia

Rosuvastatin is indicated as adjunctive therapy to other lipid-lowering treatments (e.g., LDL apheresis) or if such treatments are unavailable to reduce LDL-C, total-C, and ApoB in adult patients with homozygous familial hypercholesterolemia.

#### 5. Generalized atherosclerosis

Rosuvastatin is indicated as adjunctive therapy to diet to slow atherosclerosis progression in adult patients with the aim of reducing total-C and LDL-C.

Pediatrics

#### 1. Treatment of heterozygous familial hypercholesterolemia (HeFH)



It is indicated in 10 to 17-year-old pediatric patients for the treatment of heterozygous familial hypercholesterolemia (HeFH) as a supplement to diet to reduce Total-C, LDL-C and ApoB levels in male and female adolescents (at least one year after menarche). Drug treatment should be started after failing to achieve expected results with a diet therapy, and where laboratory tests show: LDL-C >190 mg/dL, or >160 mg/dL plus family history of early cardiovascular disease (CVD) or two or more cardiovascular risk factors.

## **SALCAL**

### **Anorexigenic Agent**

#### **Composition:**

Each tablet contains:  
Femproporex (as hydrochloride) 10 mg  
Excipients q.s.

#### **Presentations:**

30 tablet pack

#### **Indications:**

SALCAL is indicated in: obesity by overfeeding, obesity by sedentarism, obesity in the elderly. And in partial adeposidities: Madelung's disease, Dercum's disease, Barraquer –Simons' progressive lipodystrophy.

## SOMNO

### Nonbenzodiazepine Hypnotic

#### Composition:

Each coated tablet contains:  
Zolpidem hemitartrate 5 mg  
Excipients q.s.

Zolpidem hemitartrate 10 mg  
Excipientsq.s.

#### Presentations:

Packs containing 30 scored tablets of 5 or 10 mg

#### Indications:

Treatment of short-term insomnia, either sleep-onset insomnia, early morning awakening, or increased number of awakenings at night, for a 2 – 3 week therapy.

## **SOMNO XR**

### **Nonbenzodiazepine Hypnotic**

#### **Composition:**

Each coated sustained release tablet contains:

Zolpidem hemitartrate 12,5 mg

Excipients q.s.

#### **Presentations:**

Packs containing 30 coated bilayer tablets

#### **Indications:**

Somno XR is indicated for the treatment of insomnia characterized by the inability to fall asleep and/or to maintain sleep (measured as awakenings after falling asleep) for a 2 – 3 week therapy.

## **TELLMI**

### **Antihypertensive**

#### **Composition:**

Each tablet contains:

Telmisartan 40 mg

Excipients q.s

Telmisartan 80 mg

Excipients q.s

#### **Presentations:**

TELLMI 40 mg: 30 tablet pack

TELLMI 80 mg: 30 tablet pack

#### **Indications:**

Treatment of essential arterial hypertension. Prevention of cardiovascular morbidity and mortality in patients 55 years of age and older with a high risk of cardiovascular disease.

## TELLMI D

**Antihypertensive / Diuretic**

### Composition:

Each tablet of 40/12,5 contains:

Telmisartan		40 mg
Hydrochlorothiazide	12,5 mg	
Excipients q.s		

Each tablet of 80/12,5 contains:

Telmisartan		80 mg
Hydrochlorothiazide	12,5 mg	
Excipients q.s		

### Presentations:

TELLMI-D 40/12,5 mg: 30 tablet pack

TELLMI-D 80/12,5 mg: 30 tablet pack

### Indications:

Treatment of essential hypertension.

TELLMI-D, in fixed-dose combination (40 mg telmisartan / 12.5 mg hydrochlorothiazide and 80 mg telmisartan / 12.5 mg hydrochlorothiazide), is indicated in adults whose blood pressure can not be adequately controlled with telmisartan alone.

## **TIOF**

### **Antiglaucoma**

#### **Composition:**

Each ml of sterile ophthalmic solution contains:

Timolol (as maleate) 2,5 mg (0,25%)

Excipients q.s.

Timolol (as maleate) 5,0 mg (0,50%)

Excipients q.s.

#### **Presentations:**

Sealed drop bottle containing 10 ml of TIOF sterile ophthalmic solution 0.25 % and 0.5 %.

#### **Indications:**

Treatment of elevated intraocular pressure in patients with ocular hypertension, open-angle glaucoma, or aphakic glaucoma.

## **TIOF PLUS**

### **Antiglaucoma**

#### **Composition:**

Each 100 ml of sterile ophtalmic solution contains:

Dorzolamide (as Clorhidrate) 2 g

Timolol (as Maleate) 0,5 g

Excipients q.s.

#### **Presentations:**

Sealed drop bottle containing 6 mL of sterile ophtalmic solution

Sealed drop bottle containing 10 mL of sterile ophtalmic solution

#### **Indications:**

In the treatment of elevated intraocular pressure in patients with ocular hypertension, open-angle glaucoma, pseudoexfoliative glaucoma or other secondary open-angle glaucoma when concomitant therapy is approved.



## TOL 12 FORTE INYECTABLE

### Vitamins supplement

#### Composition:

Each ampoule of TOL 12 FORTE (5000) contains:  
Vitamin B1 (Thiamine hydrochloride) 200 mg  
Vitamin B6 (Pyridoxine hydrochloride) 100 mg  
Vitamin B12 (Hydroxocobalamin) 5000 mcg  
Excipients q.s.

Each ampoule of TOL 12 FORTE (10000) contains:  
Vitamin B1 (Thiamine hydrochloride) 200 mg  
Vitamin B6 (Pyridoxine hydrochloride) 100 mg  
Vitamin B12 (hydroxocobalamina) 10000 mcg  
Excipients q.s.

#### Presentations:

Pack containing 3 ampoules of 5,000 or 10,000

#### Indications:

Indicated in those cases where administration is required due to deficiency conditions and/or requiring a higher intake of complex B vitamins.

Uses:

Neuralgic and neuritis syndromes; intercostal neuralgia; lumbosciatic pain; shoulder-arm syndrome, trigeminal neuralgia; facial paralysis; neuritis, radiculitis, Herpes zoster; viral, deficiency, alcoholic, diabetic polyneuritis and polyneuritis in pregnancy. Sprains; luxations; fractures.

General stimulant in patients with anorexia, during convalescence and in geriatric patients.

## TOL 12 ORAL / TOL 12 FORTE

### Orexigenic

#### Composition:

Each 5 ml of oral solution contains:

Vitamin B1 15 mg  
Vitamin B6 15 mg  
Vitamin B12 50 mcg  
Excipients q.s.

Each capsule contains:

Vitamin B1 (Thiamine hydrochloride) 200 mg  
Vitamin B6 (Pyridoxine hydrochloride) 200 mg  
Vitamin B12 (Hydroxocobalamin acetate) 1000 mcg  
Excipients q.s.

#### Presentations:

250 ml bottle  
20 capsule pack

#### Indications:

Indicated in those cases where administration is required due to deficiency conditions and/or requiring a higher intake of complex B vitamins.

Uses:

Neuralgic and neuritis syndromes; intercostal neuralgia; lumbosciatic pain; shoulder-arm syndrome, trigeminal neuralgia; facial paralysis; neuritis, radiculitis, Herpes zoster; viral, deficiency, alcoholic, diabetic polyneuritis and polyneuritis in pregnancy. Sprains; luxations; fractures.

General stimulant in patients with anorexia, during convalescence and in geriatric patients.

## TOL TOTAL JARABE

### Vitamins supplement

#### Composition:

Each 100 ml of oral solution contains:  
Vitamin A palmitate 50000 I.U.  
Vitamin B1 hydrochloride 18,0 mg  
Vitamin B2 phosphate sodium 20,0 mg  
Vitamin B6 hydrochloride 28,0 mg  
Vitamin B12 0,10 mg  
Vitamin C 900,0 mg  
Vitamin D3 8000 I.U.  
Vitamin E acetate 196,68 mg  
Biotin 1,70 mg  
Dexpanthenol 70,0 mg  
Nicotinamide 220,0 mg  
Excipients q.s.

#### Presentations:

100 ml bottle

#### Indications:

Treatment and prevention of vitamin deficiency conditions.

## **TRAVOF SP**

### **Antiglaucoma**

#### **Composition:**

Each 100 ml of ophthalmic solution contains:

Travoprost 0,004 g

Excipients

#### **Presentations:**

Sealed drop bottle with 2,5 mL ophthalmic solution

#### **Indications:**

Travof® SP is indicated for the reduction of elevated intraocular pressure in adult patients with ocular hypertension or open-angle glaucoma.

Travof® SP is indicated for the reduction of elevated intraocular pressure in pediatric patients aged 2 months to <18 years with ocular hypertension or pediatric glaucoma.

## **TRAVOF T SP**

### **Antiglaucoma**

#### **Composition:**

Each 100 mL of ophthalmic solution contains:

Travoprost 0.004g

Timolol (as maleate) 0.5 g

Excipients q.s.

#### **Presentations:**

Sealed drop bottle with 2.5 mL ophthalmic solution

#### **Indications:**

Travof T SP is indicated for the reduction of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma, in whom treatment with a single drug only results in insufficient reduction of intraocular pressure.

## TREX / TREX FORTE

### Antibiotic

#### Composition:

Each TREX tablet contains:  
Azithromycin (as dihydrate) 500 mg  
Excipients q.s.

Each 5 ml of TREX oral suspension contains:  
Azithromycin (as dihydrate) 200 mg  
Excipients q.s.

Each 5 ml of TREX FORTE oral suspension contains:  
Azithromycin (as dihydrate) 400 mg  
Excipients q.s.

#### Presentations:

Pack with 3 and 6 tablets of 500 mg  
Bottle containing 15 mL of Suspension (200 mg/5mL)  
Bottle containing 30 mL of suspension (200 mg/5 mL)  
Trex suspension Forte: Bottle containing 20 ml of suspension (400 mg / 5 ml). A solvent to prepare the suspension and a dosing syringe for better dose adjustment and administration in children are included

#### Indications:

Azithromycin is indicated for the treatment of infections of the upper and lower respiratory tract, urinary tract, skin and soft tissues caused by sensitive microorganisms, supported by an antibiogram. Treatment of uncomplicated genital infections due to *Chlamydia trachomatis*. It is also indicated in the treatment of uncomplicated genital infections due to non-multiresistant *Neisseria gonorrhoeae*.

Clinical uses:

Infection of the lower respiratory tract, including bronchitis and pneumonia. Exacerbation of pulmonary obstructive chronic disease due to *H. Influenzae*, *M. Catarrhalis* and *S. Pneumoniae*.

Mild or moderate infection of the upper respiratory tract including laryngitis, pharyngitis, tonsillitis, sinusitis caused by *S. Pyogenes*.

Skin and soft tissue infections.

Azithromycin is indicated for the treatment of infections of the uncomplicated genitourinary tract or sexually transmitted infections (urethritis, endocervicitis) caused by *Clamydia trachomatis*. Likewise, it is indicated in genital infections caused by non-multiresistant *Neisseria gonorrhoeae* and *Ureaplasma urealyticum*.

## **TRIM 300**

### **Digestive Motility Regulator**

#### **Composition:**

Each sustained release coated tablet contains:  
Trimebutine maleate 300 mg  
Excipients q.s.

#### **Presentations:**

Packs with 10 and 30 sustained-release tablets

#### **Indications:**

TRIM 300 is used in irritable bowel syndrome: functional colonic diseases, inflammatory bowel disease, irritable colon, spastic colon and its manifestations such as intestinal spasms, colonic dysfunction, alternating diarrhea and constipation, meteorism, abdominal pain and distention, constipation, aftermath of gastrectomy.

# TRIOFENO

## Anti-Flu

### Composition:

Each coated tablet contains:

Ibuprofen: 200 mg

Pseudoephedrine Hydrochloride: 30 mg

Chlorpheniramine Maleate: 2 mg

Excipients q.s.

### Presentations:

Pack with 20 score tablets

### Indications:

TRIOFENO is indicated for the transient relief of nasal congestion, headache, and fever caused by flu states or common cold. TRIOFENO is useful in the management of nasal congestion caused by airway infections or inflammation and also in cases of sinusitis.



# TRIOVAL

## Anti-Flu

### Composition:

Each tablet contains:

Acetaminophen 500 mg

Pseudoephedrine hydrochloride 60 mg

Chlorphenamine maleate 4 mg

Excipients q.s.

Each 5 ml of oral suspension contains:

Acetaminophen 125 mg

Pseudoephedrine hydrochloride 30 mg

Chlorphenamine maleate 2 mg

Excipients q.s.

Each ml of drops solution contains:

Acetaminophen 120 mg

Pseudoephedrine hydrochloride 7,5 mg

Chlorphenamine maleate 0,75 mg

Excipients q.s.

### Presentations:

Tablets: packs with 10 and 80 tablets

Suspension: 100 ml suspension bottle

Oral drops: 15 ml drop bottle

### Indications:

Relief of the symptoms of flu and common cold.

# TRIOVAL DÍA y NOCHE

## Anti-Flu

### Composition:

#### Tablets:

Each white tablet (DAY) contains:

Acetaminophen 500 mg

Pseudoephedrine hydrochloride 60 mg

Excipients q.s.

Each blue tablet (NIGHT) contains:

Acetaminophen 500 mg

Pseudoephedrine hydrochloride 60 mg

Chlorphenamine maleate 4 mg

Excipients q.s.

#### Sachets:

Each sachet day for oral solution contains:

Acetaminophen: 500 mg

Pseudoephedrine Hydrochloride: 60 mg

Excipients q.s.

Each sachet night for oral solution contains:

Acetaminophen: 500 mg

Pseudoephedrine Hydrochloride: 60 mg

Chlorpheniramine Maleate: 4 mg

Excipients q.s.

### Presentations:

Tablets: Packs with 15 white tablets (day) and 5 blue tablets (night)

Sachets: Packs with 3 sachets

### Indications:

Relief of the symptoms of flu and common cold.

# TRUXIMA

## Monoclonal Antibody

### Composition:

Each mL of concentrate contains 10 mg of rituximab.

Each vial of 50 mL contains 500 mg of rituximab.

Each vial of 10 mL contains 100 mg of rituximab.

### Presentations:

- Truxima 500 mg concentrate for solution for infusion: Pack of 1 vial
- Truxima 100 mg concentrate for solution for infusion: Pack of 2 vials

### Indications:

The registered formulation of Rituximab is 10mg/ml concentrate for solution for intravenous (IV) infusion, and it is indicated for:

- Non-Hodgkin's Lymphoma (NHL): Rituximab is indicated for the treatment of previously untreated patients with stage III-IV follicular lymphoma (FL) along with chemotherapy or in patients who are chemo-resistant or are in their second and subsequent relapse after chemotherapy. It is also given in patients with CD20 positive large B cell NHL in combination with cyclophosphamide, doxorubicin, vincristine, prednisolone chemotherapy. Additionally, maintenance therapy is indicated for the treatment of FL individuals responding to induction therapy.
- Chronic Lymphocytic Leukaemia (CLL): Rituximab is indicated for the treatment of patients with previously untreated and relapsed CLL in combination with chemotherapy.
- Rheumatoid Arthritis (RA): Rituximab in combination with methotrexate is given to the adult patient with severe active RA who had insufficient response or intolerance to other disease-modifying anti-rheumatic drugs (DMARD) including one or more tumour necrosis factor inhibitor (TNFi) therapies.
- Granulomatosis with Polyangiitis and Microscopic Polyangiitis (GPA/MPA): In combination with glucocorticoids, Rituximab is given for the induction of remission in adult patients with severe, active GPA/MPA.

## **UNDERAN**

### **Antiseptic external use**

#### **Composition:**

Each 100 g of ointment contains:

Mupirocin 2 g (2%)

Excipients q.s.

#### **Presentations:**

Aluminum tube containing 15 g of UNDERAN.

#### **Indications:**

UNDERAN is indicated for the topical treatment of skin infections caused by susceptible pathogens: impetigo caused by *Staphylococcus aureus* and group A  $\beta$ -hemolytic *Streptococcus* (*Streptococcus pyogenes*), ecthyma, pyoderma, folliculitis, superinfected eczema, furunculosis, epidermolysis bullosa, superinfected dermatosis and infected traumatic skin lesions, burns. Prophylactic treatment for preventing bacterial infections and promoting rapid healing of lesions. As an alternative treatment for perianal streptococcal cellulitis.

# VALAX

## Antihypertensive

### Composition:

Each coated tablet contains:

Valsartan 80 mg

Excipients q.s.

Valsartan 160 mg

Excipients q.s.

### Presentations:

35 tablet pack

### Indications:

Arterial hypertension:

Indicated for the treatment of arterial hypertension. It may be used alone or concomitantly with other antihypertensive agents.

Heart Failure:

Treatment of heart failure stages II-IV (New York Heart Association) in patient on continuous treatment, for example, with diuretics, digitalics, or beta-blockers. The use of all these conventional treatments is not mandatory.

Post-myocardial infarction:

Valsartan is indicated to improve survival following myocardial infarction in clinically stable patients and with signs and symptoms or radiologic signs of left ventricular failure, or with left ventricular systolic dysfunction.

## **VALAX D**

**Antihypertensive / Diuretic**

### **Composition:**

Each coated tablet contains:

Valsartan 80 mg  
Hydrochlorotiazide 12,5 mg  
Excipients q.s.

Valsartan 160 mg  
Hydrochlorotiazide 12,5 mg  
Excipients q.s.

Valsartan 160 mg  
Hydrochlorotiazide 25 mg  
Excipients q.s.

### **Presentations:**

35 coated tablet pack

### **Indications:**

Treatment of hypertension in patients whose blood pressure is not adequately controlled with monotherapy

# VALAXAM

## Antihypertensive

### Composition:

Each coated tablet contains:

Valsartan 80 mg  
Amlodipine (as besylate) 5 mg  
Excipients q.s.

Valsartan 160 mg  
Amlodipine (as besylate) 5 mg  
Excipients q.s.

Valsartan 160 mg  
Amlodipine (as besylate) 10 mg  
Excipients q.s.

### Presentations:

35 tablet pack

### Indications:

Treatment of arterial hypertension in patients whose blood pressure does not respond to therapy with amlodipine or valsartan as monotherapy.

# VALAXAM D

**Antihypertensive / Diuretic**

## Composition:

Valaxam D 5/160/12,5: Each coated tablet contains:

Valsartan 160 mg  
Amlodipine Besylate 5 mg  
Hydrochlorotiazide 12,5 mg  
Excipients q.s.

Valaxam D 10/160/12,5: Each coated tablet contains:

Valsartan 160 mg  
Amlodipine Besylate 10 mg  
Hydrochlorotiazide 12,5 mg  
Excipients q.s.

Valaxam D 160/10/25: Each coated tablet contains:

Valsartan 160 mg  
Amlodipine Besylate 10 mg  
Hydrochlorotiazide 25 mg  
Excipients q.s.

Valaxam D 160/10/25: Each coated tablet contains:

Valsartan 160 mg  
Amlodipine Besylate 10 mg  
Hydrochlorotiazide 25 mg  
Excipients q.s.

Valaxam D 320/10/25: Each coated tablet contains:

Valsartan 320 mg  
Amlodipine Besylate 10 mg  
Hydrochlorotiazide 25 mg  
Excipients q.s.

## Presentations:

35 tablets pack

## Indications:

Treatment of essential arterial hypertension. This fixed-dose combination is not indicated for the initial treatment of hypertension.



## **VENLAX**

### **Antidepressant**

#### **Composition:**

Each tablet contains:

Venlafaxine (as hydrochloride) 50 mg

Excipients q.s.

Venlafaxine (as hydrochloride) 75 mg

Excipients q.s.

#### **Presentations:**

Packs with 30 scored tablets (50 mg and 75 mg)

#### **Indications:**

VENLAX is indicated for the treatment of all kinds of depression, alone or with comorbid anxiety. It is also indicated for the treatment of generalized anxiety disorders.

# VERTIUM

## Antivertigo

### Composition:

Each tablet contains:  
Diphenidol Hydrochloride 25 mg  
Excipients q.s.

### Presentations:

10 tablet pack  
40 tablet pack

### Indications:

Vertium is indicated in prevention and symptomatic treatment of peripheral vertigo associated with nausea and vomiting in conditions such as Meniere's disease or inner or middle ear surgery  
It is also indicated for prevention and treatment of nausea and vomiting associated with postoperative states, malignant neoplasms, labyrinthitis, and therapies with antineoplastic agents, radiotherapy and infectious diseases

## VITAYDE-C

### Vitamins supplement

#### Composition:

Each ml of drops solution contains:

Vitamin A 5000 I.U.

Vitamin C 75 mg

Vitamin D 1000 I.U.

Excipients q.s.

#### Presentations:

30 ml bottle

#### Indications:

Used to prevent and treat vitamin A, C and D deficiency in unweaned babies and children.

## XOLOF-D

### Ophthalmic Antibiotic / Corticosteroid

#### Composition:

Each 100 ml of sterile ophthalmic solution contains:

Tobramycin 0,3 g

Dexamethasone 0,1 g

Excipients q.s.

Each 100 g of sterile ophthalmic ointment contains:

Tobramycin 0,3 g

Dexamethasone 0,1 g

Excipients q.s.

#### Presentations:

Sealed drop bottle with 5 ml of sterile ophthalmic suspension

Tubes with 3.5 g of sterile ophthalmic ointment

#### Indications:

Treatment of steroid-responsive inflammatory ocular conditions accompanied by a bacterial ocular infection, or risk thereof. Used in inflammatory conditions of the palpebral and bulbar conjunctiva, cornea and anterior segment of the eye, in chronic anterior uveitis and corneal injury from chemical or thermal burns, radiation or penetration of foreign bodies.

## ZAFIN

### Analgesic

#### Composition:

Each coated tablet contains:  
Tramadol hydrochloride 37,5 mg  
Acetaminophen 325,0 mg  
Excipients q.s.

#### Presentations:

14 or 28 tablet pack

#### Indications:

Indicated for the treatment of moderate to severe, acute and chronic pain.

# ZIVAL

## Antihistamine

### Composition:

Each film coated tablet contains:  
Levocetirizine dihydrochloride 5 mg  
Excipients q.s.

### Presentations:

40 tablet pack

### Indications:

Indicated in prevention and treatment of skin and airway allergy syndromes, such as, seasonal and perennial allergic rhinitis; and for the treatment of idiopathic chronic urticaria.

# ZIVAL SOLUCIÓN

## Antihistamine

### Composition:

Each 5 ml of oral solution contains:  
Levocetirizine dihydrochloride 2,5 mg  
Excipients q.s.

Each ml (24 drops) of drops solution contains:  
Levocetirizine dihydrochloride 5 mg  
Excipients q.s.

### Presentations:

Oral solution: Bottle of 120 ml  
Oral drops: Dropper bottle of 20 ml

### Indications:

Indicated in prevention and treatment of skin and airway allergy syndromes, such as, seasonal and perennial allergic rhinitis; and for the treatment of idiopathic chronic urticaria.

# ZOMEL

## Antiulcer

### Composition:

Each enteric coated tablet contains:  
Esomeprazole (as magnesium salt) 20 mg  
Excipients q.s.

Each enteric coated tablet contains:  
Esomeprazole (as magnesium salt) 40 mg  
Excipients q.s.

Bioequivalence: This pharmaceutical product has demonstrated therapeutic equivalence.

### Presentations:

Zomel 20 mg: Package with 30 enteric-coated tablets  
Zomel 40 mg: Package with 30 enteric-coated tablets

### Indications:

Zomel contains a medicine called esomeprazole which belongs to a group of medicines called 'proton pump inhibitors'. These work by reducing the amount of acid the stomach produces.

Zomel is used to treat the following disorders:

Gastroesophageal reflux disease (GERD):

- Treatment of erosive reflux esophagitis
- Long-term preventive treatment of patients with healed esophagitis to prevent relapse
- Symptomatic treatment of gastroesophageal reflux disease (GERD).
- Use in adolescents from 12 years of age in the short-term treatment of gastroesophageal reflux.

In patients who need continuous treatment with non-steroidal anti-inflammatory drugs (NSAIDs):

- Indicated to reduce the incidence of gastric ulcers associated with continuous therapy with NSAIDs in patients over 18 years of age at risk of developing gastric ulcers. Patients at risk are considered to be those over 60 years of age and/or with a documented history of gastric ulcer.

For patients who require continuous treatment with low-dose aspirin (75-325 mg):

- Prevention of gastric and/or duodenal ulcers associated with low-dose aspirin treatment in patients at risk.

Following treatment with esomeprazole IV for maintenance of hemostasis and prevention of rebleeding of gastric and duodenal ulcers.

In combination with an appropriate antibacterial therapeutic regimen to eradicate *Helicobacter pylori* and to:

- Scarring of duodenal ulcer associated with *Helicobacter pylori* and
- Prevention of relapses of peptic ulcers associated with *Helicobacter pylori*.



## ZOMEL HP

### Antiulcer, Tritherapy

#### Composition:

Each dispersible tablet contains:  
Amoxicillin (As Trihydrate) 1 g  
Excipients q.s.

Each coated tablet contains:  
Clarithromycin 500 mg  
Excipients q.s.

Each capsule with enteric coated microgranules contains:  
Esomeprazole 20 mg  
Excipients q.s.

#### Presentations:

Package with 20 doses for 10 days treatment  
Package with 28 doses for 14 days treatment

#### Indications:

Combined therapy indicated for eradication of infections caused by *Helicobacter pylori* in patients with gastric and/or duodenal ulcer.